

08/16/00
JC862 U.S. PTO

08-17-00

A

DOCKET NO. : ISIS-4407

PATENT

JC862 U.S. PTO
09/640279
08/16/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In Re Application of:

Yogesh S. Sanghvi and Quaniai Song

Serial No.: N/A

Group Art Unit: N/A

Filing Date: Herewith

Examiner: N/A

For: PROCESSES FOR THE PREPARTION OF OLIGONUCLEOTIDES

EXPRESS MAIL LABEL NO: EM405755208US

DATE OF DEPOSIT: August 16, 2000

Box ☒ Patent Application
☐ Provisional ☐ Design

Assistant Commissioner for Patents
Washington DC 20231

PATENT APPLICATION TRANSMITTAL LETTER

Transmitted herewith for filing, please find

☒ A Utility Patent Application under 37 C.F.R. 1.53(b).

It is a continuing application, as follows:

☐ continuation ☐ divisional ☐ continuation-in-part of prior application number
____/____.

☐ A Provisional Patent Application under 37 C.F.R. 1.53(c).

☐ A Design Patent Application (submitted in duplicate).

Including the following:

☐ Provisional Application Cover Sheet.

☒ New or Revised Specification, including pages 1 to 85 containing:

- ☒ Specification
- ☒ Claims
- ☒ Abstract
- ☐ Substitute Specification, including Claims and Abstract.

☐ The present application is a continuation application of Application No. _____ filed _____. The present application includes the Specification of the parent application which has been revised in accordance with the amendments filed in the parent application. Since none of those amendments incorporate new matter into the parent application, the present revised Specification also does not include new matter.

☐ The present application is a continuation application of Application No. _____ filed _____, which in turn is a continuation-in-part of Application No. _____ filed _____. The present application includes the Specification of the parent application which has been revised in accordance with the amendments filed in the parent application. Although the amendments in the parent C-I-P application may have incorporated new matter, since those are the only revisions included in the present application, the present application includes no new matter in relation to the parent application.

☐ A copy of earlier application Serial No. _____ Filed _____, including Specification, Claims and Abstract (pages 1 - @@), to which no new matter has been added TOGETHER WITH a copy of the executed oath or declaration for such earlier application and all drawings and appendices. Such earlier application is hereby incorporated into the present application by reference.

☐ Please enter the following amendment to the Specification under the Cross-Reference to Related Applications section (or create such a section) : "This Application:

☐ is a continuation of ☐ is a divisional of ☐ claims benefit of U.S. provisional Application Serial No. _____ filed _____

☐ Signed Statement attached deleting inventor(s) named in the prior application.

- ☐ A Preliminary Amendment.
- ☐ _____ Sheets of ☐ Formal ☐ Informal Drawings.
- ☐ Petition to Accept Photographic Drawings.
- ☐ Petition Fee
- ☒ An ☐ Executed ☒ Unexecuted Declaration or Oath and Power of Attorney.
- ☒ An Associate Power of Attorney.
- ☐ An ☐ Executed ☐ Copy of Executed Assignment of the Invention to _____
- ☐ A Recordation Form Cover Sheet.
- ☐ Recordation Fee - \$40.00.
- ☐ The prior application is assigned of record to _____
- ☐ Priority is claimed under 35 U.S.C. § 119 of Patent Application No. _____
filed _____ in _____ (country).
- ☐ A Certified Copy of each of the above applications for which priority is claimed:
- ☐ is enclosed.
- ☐ has been filed in prior application Serial No. _____ filed _____.
- ☐ An ☐ Executed or ☐ Copy of Executed Earlier Statement Claiming Small Entity Status under 37 C.F.R. 1.9 and 1.27
- ☐ is enclosed.
- ☐ has been filed in prior application Serial No. _____ filed _____, said status is still proper and desired in present case.
- ☐ Diskette Containing DNA/Amino Acid Sequence Information.
- ☐ Statement to Support Submission of DNA/Amino Acid Sequence Information.
- ☐ The computer readable form in this application _____, is identical with that filed in Application Serial Number _____, filed _____. In accordance with 37 CFR 1.821(e), please use the ☐ first-filed, ☐ last-filed or ☐ only computer readable

form filed in that application as the computer readable form for the instant application. It is understood that the Patent and Trademark Office will make the necessary change in application number and filing date for the computer readable form that will be used for the instant application. A paper copy of the Sequence Listing is ☐ included in the originally-filed specification of the instant application, ☐ included in a separately filed preliminary amendment for incorporation into the specification.

- ☐ Information Disclosure Statement.
- ☐ Attached Form 1449.
- ☐ Copies of each of the references listed on the attached Form PTO-1449 are enclosed herewith.
- ☐ A copy of Petition for Extension of Time as filed in the prior case.
- ☐ Appended Material as follows: _____.
- ☒ Return Receipt Postcard (should be specifically itemized).
- ☐ Other as follows: _____

_____.

003820"62241560

FEE CALCULATION:

- ☐ Cancel in this application original claims _____ of the prior application before calculating the filing fee. (At least one original independent claim must be retained for filing purposes.)


			SMALL ENTITY		NOT SMALL ENTITY	
			RATE	FEE	RATE	FEE
PROVISIONAL APPLICATION			\$75.00	\$	\$150.00	\$
DESIGN APPLICATION			\$155.00	\$	\$310.00	\$
UTILITY APPLICATIONS BASE FEE			\$345.00	\$	\$690.00	\$ 690.00
UTILITY APPLICATION; ALL CLAIMS CALCULATED AFTER ENTRY OF ALL AMENDMENTS						
	No. Filed	No. Extra				
TOTAL CLAIMS	78 - 20 =	58	\$9 each	\$	\$18 each	\$1,044.00
INDEP. CLAIMS	4 - 3 =	1	\$39 each	\$	\$78 each	\$78.00
FIRST PRESENTATION OF MULTIPLE DEPENDENT CLAIM			\$130	\$	\$260	\$
ADDITIONAL FILING FEE				\$		\$1,122.00
TOTAL FILING FEE DUE				\$		\$1,812.00

- ☒ A Check is enclosed in the amount of \$ 1,812.00.
- ☒ The Commissioner is authorized to charge payment of the following fees and to refund any overpayment associated with this communication or during the pendency of this application to deposit account 23-3050. This sheet is provided in duplicate.
- ☐ The foregoing amount due.
- ☒ Any additional filing fees required, including fees for the presentation of extra claims under 37 C.F.R. 1.16.
- ☒ Any additional patent application processing fees under 37 C.F.R. 1.17 or 1.20(d).
- ☐ The issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance.
- ☒ The Commissioner is hereby requested to grant an extension of time for the appropriate length of time, should one be necessary, in connection with this filing or any future filing submitted to the U.S. Patent and Trademark Office in the above-identified application during the pendency of this application. The Commissioner is

further authorized to charge any fees related to any such extension of time to deposit account 23-3050. This sheet is provided in duplicate.

SHOULD ANY DEFICIENCIES APPEAR with respect to this application, including deficiencies in payment of fees, missing parts of the application or otherwise, the United States Patent and Trademark Office is respectfully requested to promptly notify the undersigned.

Date: August 16, 2000



Joseph Lucci
Registration No. 33,307

Woodcock Washburn Kurtz
Mackiewicz & Norris LLP
One Liberty Place - 46th Floor
Philadelphia PA 19103
Telephone: (215) 568-3100
Facsimile: (215) 568-3439

PROCESSES FOR THE PREPARATION OF OLIGONUCLEOTIDES**FIELD OF THE INVENTION**

The present invention is directed to methods for
5 synthesizing oligonucleotides and analogs thereof. In one
aspect of the invention the methods combine oxidation and
capping into a single step to improve the efficiency of
synthesis. The overall synthesis preferably is completed in
less time with a reduction in bulk reagents required. More
10 specific objectives and advantages of the invention will
hereinafter be made clear or become apparent to those
skilled in the art during the course of explanation of
preferred embodiments of the invention.

BACKGROUND OF THE INVENTION

15 Modified oligonucleotides are of great value in
molecular biological research and in applications such as
anti-viral therapy. Modified oligonucleotides which can
block RNA translation, and are nuclease resistant, are
useful as antisense reagents. In addition to oligonucleo-
20 tides that have phosphodiester internucleotide linkages,
sulfurized oligonucleotides which contain, for example,
phosphorothioate linkages are also of interest in these
areas. Because of their chirality (Rp and Sp)
phosphorothioate containing oligonucleotides are useful in
25 determining stereochemical pathways of certain enzymes which
recognize nucleic acids.

It is well known that most of the bodily states in
multicellular organisms, including most disease states, are

effected by proteins. Such proteins, either acting directly or through their enzymatic or other function, contribute in major proportion to many diseases and regulatory functions in animals and humans. For disease states, classical
5 therapeutics has generally focused upon interactions with such proteins in efforts to moderate their disease-causing or disease-potentiating functions. In newer therapeutic approaches, modulation of the actual production of such proteins is desired. By interfering with the production of
10 proteins, the maximum therapeutic effect may be obtained with minimal side effects. It is therefore a general object of such therapeutic approaches to interfere with or otherwise modulate gene expression, which would lead to undesired protein formation.

15 One method for inhibiting specific gene expression is with the use of oligonucleotides, especially oligonucleotides which are complementary to a specific target messenger RNA (mRNA) sequence. Several oligonucleotides are currently undergoing clinical trials for such use. Phosphorothioate
20 oligonucleotides are presently being used as such antisense agents in human clinical trials for various disease states, including use as antiviral agents.

Transcription factors interact with double-stranded DNA during regulation of transcription. Oligonucleotides can
25 serve as competitive inhibitors of transcription factors to modulate their action. Several recent reports describe such interactions (see Bielinska, A., et. al., *Science*, 1990, 250, 997-1000; and Wu, H., et. al., *Gene*, 1990, 89, 203-209).

30 In addition to such use as both indirect and direct regulators of proteins, oligonucleotides and their analogs also have found use in diagnostic tests. Such diagnostic tests can be performed using biological fluids, tissues, intact cells or isolated cellular components. As with gene

expression inhibition, diagnostic applications utilize the ability of oligonucleotides and their analogs to hybridize with a complementary strand of nucleic acid. Hybridization is the sequence specific hydrogen bonding of oligomeric
5 compounds via Watson-Crick and/or Hoogsteen base pairs to RNA or DNA. The bases of such base pairs are said to be complementary to one another.

Oligonucleotides and their analogs are also widely used as research reagents. They are useful for understanding the
10 function of many other biological molecules as well as in the preparation of other biological molecules. For example, the use of oligonucleotides and their analogs as primers in PCR reactions has given rise to an expanding commercial industry. PCR has become a mainstay of commercial and
15 research laboratories, and applications of PCR have multiplied. For example, PCR technology now finds use in the fields of forensics, paleontology, evolutionary studies and genetic counseling. Commercialization has led to the development of kits which assist non-molecular biology-
20 trained personnel in applying PCR. Oligonucleotides and their analogs, both natural and synthetic, are employed as primers in such PCR technology.

Oligonucleotides and their analogs are also used in other laboratory procedures. Several of these uses are
25 described in common laboratory manuals such as *Molecular Cloning, A Laboratory Manual*, Second Ed., J. Sambrook, et al., Eds., Cold Spring Harbor Laboratory Press, 1989; and *Current Protocols In Molecular Biology*, F. M. Ausubel, et al., Eds., Current Publications, 1993. Such uses include as
30 synthetic oligonucleotide probes, in screening expression libraries with antibodies and oligomeric compounds, DNA sequencing, *in vitro* amplification of DNA by the polymerase chain reaction, and in site-directed mutagenesis of cloned DNA. See Book 2 of *Molecular Cloning, A Laboratory Manual*,

supra. See also "DNA-protein interactions and The Polymerase Chain Reaction" in Vol. 2 of *Current Protocols In Molecular Biology*, *supra*.

Oligonucleotides and their analogs can be synthesized
5 to have customized properties that can be tailored for
desired uses. Thus a number of chemical modifications have
been introduced into oligomers to increase their usefulness
in diagnostics, as research reagents and as therapeutic
entities. Such modifications include those designed to
10 increase binding to a target strand (i.e. increase their
melting temperatures, T_m), to assist in identification of
the oligonucleotide or an oligonucleotide-target complex, to
increase cell penetration, to stabilize against nucleases
and other enzymes that degrade or interfere with the
15 structure or activity of the oligonucleotides and their
analogs, to provide a mode of disruption (terminating event)
once sequence-specifically bound to a target, and to improve
the pharmacokinetic properties of the oligonucleotide.

The synthesis of oligonucleotides has classically
20 involved obtaining a desired product on a small scale. The
synthesis of oligonucleotides has more recently evolved to
the point that routine syntheses are being performed on
kilogram scale. Moving forward the next step is the
synthesis of oligonucleotides and analogs on ton scale to
25 supply large quantities to meet demands for ongoing
pharmaceutical sales and clinical trials. The evolution of
oligonucleotide synthetic techniques toward larger scale is
requiring a reevaluation of each aspect of the synthetic
process. There is an ongoing need in the art of oligomer
30 synthesis to improve the efficiency of synthesis.

The chemical literature discloses numerous protocols
for coupling nucleosides through phosphorous-containing
covalent linkages to produce oligonucleotides of defined
sequence. One of the most routinely used protocols is the

phosphoramidite protocol (see, e.g., Advances in the Synthesis of Oligonucleotides by the Phosphoramidite Approach, Beaucage, S.L.; Iyer, R.P., *Tetrahedron*, **1992**, *48*, 2223-2311 and references cited therein; and The synthesis of Modified Oligonucleotides by the Phosphoramidite Approach and their applications, Beaucage, S.L.; Iyer, R.P., *Tetrahedron*, **1993**, *49*, 6123-6194 and references cited therein), wherein a nucleoside or oligonucleotide having a free hydroxyl group is reacted with a protected phosphoramidite monomer in the presence of a weak acid to form a phosphite-linked structure. Oxidation of the phosphite linkage with a suitable reagent effects conversion of a P^{III} internucleoside linkage to a P^V internucleoside linkage. For the purpose of this application, such reagents include oxygen transfer reagents and sulfur transfer reagents. Subsequent hydrolysis of the cyanoethyl group yields the desired phosphodiester or phosphorothioate linkage.

Phosphoramidites are commercially available from a variety of commercial sources (included are: Glen Research, Sterling, Virginia; Amersham Pharmacia Biotech Inc., Piscataway, New Jersey; Cruachem Inc., Aston, Pennsylvania; Chemgenes Corporation, Waltham, Massachusetts; Proligo LLC, Boulder, Colorado; PE Biosystems, Foster City California; Beckman Coulter Inc., Fullerton, California).

An efficient sulfur transfer reagent is important to the success of obtaining high quality phosphorothioate product with a low percentage of phosphate linkages. A number of sulfur transfer reagents have been reported, including Beaucage reagent, 3-ethoxy-1,2,4-dithiozoline-5-one (EDITH), 1,2,4-dithiozoline-3,5-dione, 3-methyl-ethoxy-1,2,4-dithiozoline-5-one (MEDITH), phenylacetyl disulfide (PADS), tetraethylthiuram disulfide (TETD) and others. The cost of sulfur transfer reagents and their efficiency has to be taken into account when performing large-scale

manufacturing. Beaucage reagent is expensive (about \$5.00 per gram) and is being replaced by other cheaper reagents like PADS. PADS was first introduced by van Boom but did not show adequate sulfur transfer efficiency under the
5 original conditions, which did not use a base during the sulfurization step (Roclen et al., *Recl. Trav. Chim. Pays-Bas*, 1991, 110, 325-331).

The use of elemental sulfur in oligonucleotide synthesis presents problems and is not suitable for automation because
10 of sulfur's insolubility in most organic solvents.

Furthermore, carbon disulfide, a preferred source of sulfur, has undesirable volatility and an undesirably low flash point. Unwanted side products are often observed with the use of dibenzoyl tetrasulfide. Beaucage reagent, while a
15 relatively efficient sulfurization reagent, is difficult to synthesize and not particularly stable. Furthermore, use of Beaucage reagent forms a secondary reaction product which is a potent oxidizing agent. (R.P. Iyer et al., *J. Am. Chem. Soc.* 112, pp. 1253-1254 (1990); R. P. Iyer et al., *J. Org.*
20 *Chem.* 55, 4693-4699 (1990)). This can further lead to unwanted side products which can be difficult to separate from the desired reaction product. Tetraethylthiuram disulfide while relatively inexpensive and stable, has a sulfurization reaction rate which can be undesirable slow.

25 A method of preparing phosphorothioate oligonucleotides using tetraethylthiuram disulfide is disclosed in United States Patent Number 5,166,387. Although the use of tetraethylthiuram disulfide as a sulfur transfer reagent has been described since 1992, it has not been used for
30 commercial scale production of phosphorothioate oligonucleotides. Beaucage reagent and phenylacetyl disulfide (PADS) are the only reagents that have been used commercially with good results. Our experiments with tetraethylthiuram disulfide indicate that there is a


009489 "6234360

significant amount of side product formation (see Example 8) along with the desired phosphorothioate product. Although we do not wish to be bound by theory it is believed that tetraethylthiuram disulfide is an overly stable sulfur transfer reagent that is difficult to dissociate after reaction with the P^{III} species.

Thus, there remains a need for improved methods and reagents for preparing oligomeric compounds. The present invention is directed to these, as well as other, important ends.

SUMMARY OF THE INVENTION

The present invention provides methods of preparing oligomeric compounds having at least one moiety of formula:



wherein:

X_2 is O or S;

X_1 is Pg-O-, Pg-S-, C_1 - C_{10} straight or branched chain alkyl, $CH_3(CH_2)_{nn}$ -O-, R_2R_3N - or a group remaining from coupling a chiral auxiliary;

nn is from 0 to 10;

Pg is CH_3 , $-CH_2CH_2CN$, $-C(CH_3)(CH_3)-CCl_3$, $-CH_2-CCl_3$, $-CH_2CH=CH_2$, $CH_2CH_2SiCH_3$, 2-yl-ethyl phenylsulfonate, δ -cyano-butenyl, cyano *p*-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-

trifluoroacetyl ethyl, acetoxo phenoxy ethyl, or a blocking group;

each R_2 and R_3 is, independently, hydrogen, C_1 - C_{10} alkyl, cycloalkyl or aryl;


- 5 or optionally, R_2 and R_3 , together with the nitrogen atom to which they are attached form a cyclic moiety that may include an additional heteroatom selected from O, S and N;

each B_x is, independently, a heterocyclic base moiety;

10 and

each R_1 is, independently, H, a blocked hydroxyl group, or a sugar substituent group; comprising the steps of:

- (a) providing a 5'-O-protected compound of the
15 formula:




wherein:

T₁ is a hydroxyl protecting group; and

20 T₂ is a covalent attachment to a support media, a nucleoside bound to a support media, a nucleotide, an oligonucleoside or an oligonucleotide;

(b) treating said 5'-O-protected compound with a deprotecting reagent for a time and under conditions effective to form a 5'-O-deprotected compound;

- 25 (c) coupling said 5'-O-deprotected compound with an activated phosphorus composition of the formula:



wherein:

T₃ is a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleoside or an oligonucleotide;

5 R₄ is N(L₁)L₂,


each L₁ and L₂ is, independently, C₁₋₆ straight or branched alkyl, or a C₅₋₇ cyclic aliphatic ring system;

or L₁ and L₂ are joined together to form a 4- to 13-membered heterocyclic ring system including the nitrogen
10 atom to which L₁ and L₂ are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S; and

R₅ is X₁;

or R₄ and R₅ together with the phosphorus atom to which
15 R₄ and R₅ are attached form a chiral auxiliary;

for a time and under conditions effective to form an extended compound having the formula:



(d) treating said extended compound with a mixture comprising an oxidizing reagent and a capping reagent for a time and under conditions effective to form said oligomeric compound.

5 The methods can include further treatment of the oligomeric compound with a reagent for a time and under conditions effective to remove the blocking groups, thereby forming a deblocked oligomeric compound. This reagent can also be effective to cleave the oligomeric compound from the
10 support media. One such reagent is aqueous ammonium hydroxide. Alternatively, the deblocking can be effected with one reagent, and a further reagent can effect cleavage from the support media. Also included is treatment with a deprotecting reagent for a time and under conditions
15 effective to deprotect the T₃ hydroxyl protecting group.

In one embodiment of the invention, the mixture comprises from 0.02M to 0.2M oxidizing reagent with a preferred range is from 0.1M to 0.2M.

The oxidizing reagent can be one that transfers an
20 oxygen atom. Effective oxidizing reagents for oxygen transfer are iodine, *m*-chloroperbenzoic acid, iodobenzene diacetate, tetra-*n*-butylammonium periodate, *tert*-butyl hydroperoxide, di-*tert*-butyl hydroperoxide, cumene hydroperoxide, hydrogen peroxide; bis-trimethylsilyl
25 peroxide; dinitrogen tetroxide, oxone, molecular oxygen, (1*S*)-(+)-(10-camphorsulfonyl)oxaziridine or a peracid with iodine, *m*-chloroperbenzoic acid, iodobenzene diacetate, *tert*-butyl hydroperoxide, di-*tert*-butyl hydroperoxide, hydrogen peroxide; oxone, molecular oxygen or a peracid being
30 preferred.

The oxidizing reagent can also be one that transfers a sulfur atom. Effective oxidizing reagents for sulfur transfer are 3-amino-1,2,4-dithiazole-5-thione; 3-ethoxy-1,2,4-dithiazoline-5-one; 1,2,4-dithiazolidine-3,5-dione; 3-

methyl-1,2,4-dithiazolin-5-one; or dimethylthiuram disulfide with dimethylthiuram disulfide being preferred.

In one embodiment, the capping reagent is prepared by mixing equal volumes of two components prior to the capping
5 step. The first component is a mixture of an acylating agent such as N-methylimidazole or 4-dimethylaminopyridine and a base such as pyridine or 2,6-lutidine in acetonitrile or tetrahydrofuran. The second component is a solution of an
10 anhydride such as acetic anhydride, chloroacetic anhydride, t-butylphenoxyacetic anhydride in acetonitrile or tetrahydrofuran.

One preferred capping reagent comprises about equal volumes of a first component containing from 5% to about 25% N-methylimidazole, from about 20% to about 50% pyridine and
15 from about 20% to about 50% acetonitrile, added to a second component containing from about 20% to about 50% acetic anhydride in acetonitrile.

Another preferred capping reagent comprises about equal volumes of a first component containing about 5% to about 25%
20 4-dimethylaminopyridine, from about 20% to about 50% 2,6-lutidine in acetonitrile, added to a second component containing from about 20% to about 50% acetic anhydride in acetonitrile.

The mixture of oxidizing reagent and capping reagent can
25 comprise, for example, dimethylthiuram disulfide, acetic anhydride, acetonitrile, N-methyl imidazole, or pyridine. A preferred mixture comprises from about 0.05M to 0.2M dimethylthiuram disulfide, about 10% acetic anhydride, about 10% N-methyl imidazole and about 15% pyridine in a suitable
30 solvent. Suitable solvents include acetonitrile, toluene, ethyl acetate, tetrahydrofuran, dichloromethane, dichloroethane, dioxane, dimethylacetamide and dimethylformamide.


In another embodiment of the invention the coupling of the 5'-O-deprotected compound with the activated phosphorus composition is performed in the presence of an activating agent that renders the phosphorous atom more susceptible to nucleophilic attack. Preferred activating agents include 1-H-tetrazole and 4,5-dicyanoimidazole.

Exemplary cyclic moieties according to the invention include morpholino or phthalimido moieties.

Each L_1 and L_2 can be, independently, C_{1-6} alkyl with isopropyl being a preferred alkyl group for L_1 and L_2 . L_1 and L_2 can also be joined together to form a heterocyclic ring system including the nitrogen atom to which said L_1 and L_2 are attached, the ring system optionally includes at least one additional heteroatom selected from O, N and S. A preferred heterocyclic ring system is morpholino.

Representative substituent groups according to the invention include, C_1-C_{20} alkyl, C_2-C_{20} alkenyl, C_2-C_{20} alkynyl, C_5-C_{20} aryl, O-alkyl, O-alkenyl, O-alkynyl, O-aryl, O-aralkyl, O-alkylamino, O-alkylaminoalkyl (O-alkyl-N(H)alkyl), O-alkylaminodialkyl (O-alkyl-N-(alkyl)₂), O-alkylalkoxy (O-alkyl-O-alkyl), O-alkyl-(N-imidazole), thiol, S-alkyl, S-alkenyl, S-alkynyl, NH-alkyl, NH-alkenyl, NH-alkynyl, N-dialkyl, S-aryl, NH-aryl, S-aralkyl, NH-aralkyl, N-phthalimido, halogen keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, N-imidazole, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocycle, carbocycle, polyamine, polyamide, polyalkylene glycol, or polyether;

Alternatively, one or more substituent groups has one of formula I or II:




wherein:

Z₀ is O, S or NH;

J is a single bond, O or C(=O);

5 E is C₁-C₁₀ alkyl, N(R₁)(R₂), N(R₁)(R₅), N=C(R₁)(R₂),
N=C(R₁)(R₅) or has one of formula III or IV;



each R₆, R₇, R₈, R₉ and R₁₀ is, independently, hydrogen,
C(O)R₁₁, substituted or unsubstituted C₁-C₁₀ alkyl, substituted
10 or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted
C₂-C₁₀ alkynyl, alkylsulfonyl, arylsulfonyl, a chemical
functional group or a conjugate group, wherein the
substituent groups are selected from hydroxyl, amino, alkoxy,
carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen,
15 alkyl, aryl, alkenyl and alkynyl;

or optionally, R₇ and R₈, together form a phthalimido
moiety with the nitrogen atom to which they are attached;

or optionally, R₉ and R₁₀, together form a phthalimido
moiety with the nitrogen atom to which they are attached;

20 each R₁₁ is, independently, substituted or unsubstituted
C₁-C₁₀ alkyl, trifluoromethyl, cyanoethoxy, methoxy, ethoxy,
t-butoxy, allyloxy, 9-fluorenylmethoxy, 2-(trimethylsilyl)-
ethoxy, 2,2,2-trichloroethoxy, benzyloxy, butyryl, iso-
butyryl, phenyl or aryl;

R_5 is T-L,

T is a bond or a linking moiety;

L is a chemical functional group, a conjugate group or a support media;

5 each R_1 and R_2 is, independently, H, a nitrogen protecting group, substituted or unsubstituted C_1 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} alkenyl, substituted or unsubstituted C_2 - C_{10} alkynyl, wherein said substitution is OR_3 , SR_3 , NH_3^+ , $N(R_3)(R_4)$, guanidino or acyl where said acyl is
10 an acid amide or an ester;

or R_1 and R_2 , together, are a nitrogen protecting group or are joined in a ring structure that optionally includes an additional heteroatom selected from N and O;

or R_1 , T and L, together, are a chemical functional
15 group;

each R_3 and R_4 is, independently, H, C_1 - C_{10} alkyl, a nitrogen protecting group, or R_3 and R_4 , together, are a nitrogen protecting group;

or R_3 and R_4 are joined in a ring structure that
20 optionally includes an additional heteroatom selected from N and O;

Z_4 is OX, SX, or $N(X)_2$;

each X is, independently, H, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $C(=NH)N(H)R_5$, $C(=O)N(H)R_5$ or $OC(=O)N(H)R_5$;

25 R_5 is H or C_1 - C_8 alkyl;

Z_1 , Z_2 and Z_3 comprise a ring system having from about 4 to about 7 carbon atoms or having from about 3 to about 6 carbon atoms and 1 or 2 hetero atoms wherein said hetero atoms are selected from oxygen, nitrogen and sulfur and
30 wherein said ring system is aliphatic, unsaturated aliphatic, aromatic, or saturated or unsaturated heterocyclic;

Z_5 is alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl

Pub. No. 4,204,360

having 2 to about 10 carbon atoms, aryl having 6 to about 14 carbon atoms, $N(R_1)(R_2)OR_1$, halo, SR_1 or CN;

each q_1 is, independently, an integer from 1 to 10;

each q_2 is, independently, 0 or 1;

5 q_3 is 0 or an integer from 1 to 10;

q_4 is an integer from 1 to 10;

q_5 is from 0, 1 or 2; and

provided that when q_3 is 0, q_4 is greater than 1.

In one embodiment of the present invention X_1 is Pg-O-,
10 Pg-S-, -CH₃, CH₃-O-, morpholino or -NR₂R₃ where each R₂ and R₃ is, independently, hydrogen or C₁-C₁₀ alkyl. Where Pg is CH₂CH₂CN, diphenylsilylethyl, δ -cyanobutenyl, cyano *p*-xylyl, methyl-N-trifluoroacetyl ethyl or acetoxy phenoxy ethyl.

Representative heterocyclic bases include adenine, N⁶-
15 benzoyladenine, cytosine, N⁴-benzoylcytosine, 5-methylcytosine, N⁴-benzoyl-5-methylcytosine, thymine, uracil, guanine, N²-isobutyrylguanine and 2-aminoadenine, as well as protected (or "blocked") forms thereof.

In certain embodiments of the present invention, the
20 support media bound nucleoside, nucleotide, oligonucleoside or oligonucleotide is blocked at reactive sites. In one embodiment the blocking groups are acid stable. In another embodiment the blocking groups are base labile.


Deprotecting reagents can be acidic, neutral or basic.
25 Preferred deprotecting reagents are dichloroacetic acid, trichloroacetic acid, zinc bromide, AlCl₃, TiCl₄, (Et)AlCl, (i-Bu)₂AlCl, ceric ammonium nitrate, 1,1,1,3,3,3-hexafluoro-2-propanol, and diethyloxomalonate. A particularly preferred deprotecting reagent is 2-5% dichloroacetic acid in
30 dichloromethane or dichloroethane. In a further embodiment the deprotecting reagent is a fluoride moiety where a preferred fluoride moiety is BF₃-etherate.

The oligomeric compounds of the invention typically comprises from 5 to about 50 nucleosides, with from 8 to

003430" 04204360

about 30 nucleosides being preferred and from 15 to about 25 nucleosides being more preferred.

The present invention also provides methods for the preparation of oligomeric compounds having at least one moiety of formula:



wherein:

X_1 is Pg-O-, Pg-S-, C_1 - C_{10} straight or branched chain alkyl, $CH_3(CH_2)_{nn}$ -O-, R_2R_3N - or a group remaining from coupling
10 a chiral auxiliary;

nn is from 0 to 10;

Pg is CH_3 , $-CH_2CH_2CN$, $-C(CH_3)(CH_3)-CCl_3$, $-CH_2-CCl_3$, $-CH_2CH=CH_2$, $CH_2CH_2SiCH_3$, 2-yl-ethyl phenylsulfonate, δ -cyanobutenyl, cyano *p*-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-
15 trifluoroacetyl ethyl, acetoxy phenoxy ethyl, or a blocking group;

each R_2 and R_3 is, independently, hydrogen, C_1 - C_{10} alkyl, cycloalkyl or aryl;


20 or optionally, R_2 and R_3 , together with the nitrogen atom to which they are attached form a cyclic moiety that may include an additional heteroatom selected from O, S and N;

each Bx is, independently, a heterocyclic base moiety or a blocked heterocyclic base moiety; and

each R_1 is, independently, H, a blocked hydroxyl group, a sugar substituent group or a blocked sugar substituent group;

comprising the steps of:

- 5 (a) providing a 5'-O-protected compound of the formula:




wherein:

T_1 is a hydroxyl protecting group; and

T_2 is a covalent attachment to a support media, or a
10 support media bound nucleoside, nucleotide, oligonucleoside or oligonucleotide;

(b) treating said 5'-O-protected compound with a deprotecting reagent for a time and under conditions effective to form a 5'-O-deprotected compound;

- 15 (c) coupling said 5'-O-deprotected compound with an activated phosphorus composition of the formula:



wherein:

T_3 is a hydroxyl protecting group, a nucleoside, a
20 nucleotide, an oligonucleoside or an oligonucleotide;

R_4 is $N(L_1)L_2$;


each L_1 and L_2 is, independently, C_{1-6} straight or branched alkyl, or a C_{5-7} cyclic aliphatic ring system;

or L_1 and L_2 are joined together to form a 4- to 13-membered heterocyclic ring system including the nitrogen atom to which L_1 and L_2 are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S; and


R_5 is X_1 ;

or R_4 and R_5 together with the phosphorus atom to which R_4 and R_5 are attached form a chiral auxiliary;

for a time and under conditions effective to form an extended compound having the formula:




(d) treating said extended compound with dimethylthiuram disulfide in a solvent thereby forming a sulfurized compound having the formula:



(e) treating said sulfurized compound with a capping reagent for a time and under conditions effective to form said oligomeric compound.

- 5 In one embodiment the dimethylthiuram disulfide is from about 0.02M to about 0.2M in said solvent. In a preferred embodiment the dimethylthiuram disulfide is from about 0.1M to about 0.2M in said solvent.

The present invention also provides methods of preparing
10 oligomeric compounds having at least one moiety having one of the formulas:



wherein:

X_2 is O or S;

X_1 is $Pg-O-$, $Pg-S-$, C_1-C_{10} straight or branched chain alkyl, $CH_3(CH_2)_{nn}-O-$, R_2R_3N- or a group remaining from coupling a chiral auxiliary;

nn is from 0 to 10;

5 Pg is CH_3 , $-CH_2CH_2CN$, $-C(CH_3)(CH_3)-CCl_3$, $-CH_2-CCl_3$, $-CH_2CH=CH_2$, $CH_2CH_2SiCH_3$, 2-yl-ethyl phenylsulfonate, δ -cyano-butenyl, cyano *p*-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-trifluoroacetyl ethyl, acetoxy phenoxy ethyl, or a blocking
10 group;

each R_2 and R_3 is, independently, hydrogen, C_1-C_{10} alkyl, cycloalkyl or aryl;


or optionally, R_2 and R_3 , together with the nitrogen atom to which they are attached form a cyclic moiety that may
15 include an additional heteroatom selected from O, S and N;

each Bx is, independently, a heterocyclic base moiety or a blocked heterocyclic base moiety; and

each R_1 is, independently, H, a blocked hydroxyl group, a sugar substituent group or a blocked sugar substituent
20 group;

comprising the steps of:

(a) providing a 5'-O-protected compound having one of the formulas:



25 or


wherein:

T_1 is a hydroxyl protecting group; and

T_2 is a covalent attachment to a support media, or a support media bound nucleoside, nucleotide, oligonucleoside
30 or oligonucleotide;

(b) treating said 5'-O-protected compound with a deprotecting reagent for a time and under conditions effective to form a 5'-O-deprotected compound;

(c) coupling said 5'-O-deprotected compound with an activated phosphorus composition of the formula:



wherein:

T₃ is a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleoside or an oligonucleotide;

10 R₄ is N(L₁)L₂;

each L₁ and L₂ is, independently, C₁₋₆ straight or branched alkyl, or a C₅₋₇ cyclic aliphatic ring system;


or L₁ and L₂ are joined together to form a 4- to 13-membered heterocyclic ring system including the nitrogen atom
15 to which L₁ and L₂ are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S; and

R₅ is X₁;

or R₄ and R₅ together with the phosphorus atom to which
20 R₄ and R₅ are attached form a chiral auxiliary;

for a time and under conditions effective to form an extended compound having one of the formulas:

003760-04241350




and

(d) treating said extended compound with a mixture comprising an oxidizing reagent and a capping reagent for a time and under conditions effective to form said oligomeric compound.

DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention is directed to methods for synthesizing oligonucleotides and analogs thereof having at least one moiety of one of the formulas:



wherein:

X_2 is O or S;

X_1 is Pg-O-, Pg-S-, C_1 - C_{10} straight or branched chain alkyl, $CH_3(CH_2)_{nn}$ -O-, R_2R_3N - or a group remaining from coupling a chiral auxiliary;

nn is from 0 to 10;

5 Pg is CH_3 , $-CH_2CH_2CN$, $-C(CH_3)(CH_3)-CCl_3$, $-CH_2-CCl_3$, $-CH_2CH=CH_2$, $CH_2CH_2SiCH_3$, 2-yl-ethyl phenylsulfonate, δ -cyanobutenyl, cyano *p*-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-trifluoroacetyl ethyl, acetoxy phenoxy ethyl, or a blocking
10 group;

each R_2 and R_3 is, independently, hydrogen, C_1 - C_{10} alkyl, cycloalkyl or aryl;

or optionally, R_2 and R_3 , together with the nitrogen atom to which they are attached form a cyclic moiety that may
15 include an additional heteroatom selected from O, S and N;

each Bx is, independently, a heterocyclic base moiety or a blocked heterocyclic base moiety; and

each R_1 is, independently, H, a blocked hydroxyl group, a sugar substituent group, or a blocked substituent group;

20 The present methods can further include deblocking, deprotecting and cleaving the resulting oligomeric compound. The most common deblocked forms of X_1 include -OH, -SH, $-N(R_2)(R_3)$, alkyl and alkoxy groups. Purification can be performed at various stages, but is routinely performed with
25 the terminal protecting group attached such as a trityl on purification. including protected or deprotected.

The most widely used method for the large scale synthesis of oligomeric compounds uses phosphoramidite chemistry on a support media. In general the method requires
30 4 separate and distinct steps per cycle: detritylation, coupling, oxidation and capping. Each step requires automated synthesis equipment time and significant quantities of reagents. In one aspect of the present invention, a method is disclosed that employs a mixture comprising an

oxidizing reagent and a capping reagent to simultaneously effect oxidation of internucleoside linkages and capping of unreacted hydroxyl groups. This combined oxidation and capping step is amenable to otherwise standard synthetic methods for the synthesis of oligomeric compounds.

In general, a synthon bound to a support media, e.g. a nucleoside or nucleosidic oligomer, is extended by addition of a further synthon using standard chemistries to form a phosphite or other P^{III} intermediate linkage. Treatment of this P^{III} intermediate linkage in a single step with a mixture containing an oxidizing reagent and a capping reagent will give an extended compound having the desired P^V oxidized linkage with unreacted hydroxyl groups capped. Advantages of the present methods can include a faster synthesis time and a reduction in cost due in part to the deletion of one of the reagent consuming steps and associated wash and rinse cycles. A further advantage can be gained by selecting an inexpensive oxidizing reagent or one that is easily prepared from inexpensive reagents.

Preferred mixtures of oxidizing and capping reagents include those that are stable and include components that are mutually soluble with each other. Oxidizing reagents amenable to the present invention should be soluble in the solution comprising the oxidizing reagent at concentrations of 0.02M, or greater, preferred about 0.1M or more, more preferred from 0.1 to 0.2M. A preferred mixture is an oxidizing reagent dissolved in cap A (20% acetic anhydride in acetonitrile) mixed with an equal volume of cap B (*N*-methylimidazole-pyridine-acetonitrile, 2:3:5, v/v/v). Oxidizing reagents that are soluble and stable with an equal volume mixture of cap A and cap B are preferred within the scope of the present invention.

Oxidizing reagents that are effective to transfer an oxygen atom (thereby converting a P^{III} linkage to a P^V

linkage) include without limitation *m*-chloroperbenzoic acid; iodobenzene diacetate, tetra-*n*-butylammonium periodate; *tert*-butyl hydroperoxide; di-*tert*-butyl hydroperoxide; cumene hydroperoxide; hydrogen peroxide; bis-trimethylsilyl

5 peroxide; and catalytic amounts of trimethylsilyl triflate; dinitrogen tetroxide and molecular oxygen in the presence of 2,2'-azobis(2-methylpropionitrile) under thermal or photo-chemical conditions; and (1*S*)-(+)-(10-camphorsulfonyl)-oxaziridine; iodine/tetrahydrofuran/water/pyridine; hydrogen

10 peroxide/water; *tert*-butyl hydroperoxide; and a peracid like *m*-chloroperbenzoic acid (see review article Beaucage *et al.*, *Current Protocols in Nucleic Acid Chemistry*, 2000, 3.3.1-3.3.20). In the case of oxidation to a sulfur species (sulfurization), the reaction is generally performed under

15 anhydrous conditions with an exclusion of air, e.g. oxygen. In the case of oxidation the reaction can be performed under aqueous conditions.

Oxidizing reagents that transfer a sulfur atom (sulfurizing reagents) are used to form phosphorothioate or

20 other sulfurized internucleoside linkages such as, for example, phosphorodithioate internucleoside linkages. Sulfurizing reagents amenable to the present invention include those that are partially or completely soluble in a selected capping reagent or reagents. In addition the

25 sulfurizing reagent should be compatible e.g. stable and non-reactive with the capping reagents. Preferred sulfurizing reagents are commercially available in bulk for considerably less cost than most traditional sulfurizing agents that are currently in use. Alternatively, a sulfurizing reagent is

30 selected because of its ease of synthesis from inexpensive bulk reagents.

One important criteria for a preferred sulfurizing reagent is its ability incorporate sulfur and exclude incorporation of oxygen. Analysis of an oxidized oligomer,

005480" 02204960

using ^{31}P NMR, will give the percentages of sulfurized and oxygenized internucleoside linkages.

Preferred sulfurized linkages include those that are prepared by methods known in the art to give chirally enhanced or chirally pure sulfurized linkages for those linkages that are not achiral. Preferred sulfurized linkages that are prepared by the present methods include:

- phosphorothioate $(-\text{O}-\text{P}(\text{S})(\text{O})-\text{O}-)$;
- phosphorodithioate $(-\text{O}-\text{P}(\text{S})(\text{S})-\text{O}-)$;
- 10 phosphorthioamidate $(-\text{O}-\text{P}(\text{S})(\text{NJ})-\text{O}-)$;
- phosphonothioate $(-\text{O}-\text{P}(\text{J})(\text{S})-\text{O}-)$;
- boranothiophosphate $(-\text{O}-\text{P}(\text{S})(\text{BJ}_3)-\text{J}-)$;

wherein "J" denotes a substituent group which is commonly hydrogen or an alkyl group, but which can be a more complicated group that varies from one type of linkage to another but is well known to the art skilled.

Representative United States Patents that teach the preparation of the above phosphorus atom containing linkages include, but are not limited to, U.S. Patents Nos. 3,687,808; 20 4,469,863; 4,476,301; 5,023,243; 5,166,387; 5,177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,625,050; and 25 5,697,248, certain of which are commonly owned by the assignee of this application, and each of which is herein incorporated by reference.


Positional modifications, also known in the art, involve the linking of nucleosides in a non-naturally occurring motif. As used herein the term "positional modification" is 30 meant to include without limitation 2',5'-internucleoside linkages. Combining modifications e.g. using modified chemistries and positional modifications of selected internucleoside linkages is also amenable to the present

invention where for example a 2',5'-phosphoramidate internucleoside linkage is employed. The 2'-5'-linkage has been used at the termini of oligomeric compounds to enhance the nuclease resistance (as described in U.S. Application 5 Serial No. 09/435,806, filed November 8, 1999).

A representative list of substituent groups amenable to the present invention include C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, C₅-C₂₀ aryl, O-alkyl, O-alkenyl, O-alkynyl, O-alkylamino, (O-alkyl-N(H)alkyl), O-alkylaminodialkyl (O-alkyl-N-(alkyl)₂), O-alkylalkoxy (O-alkyl-O-alkyl), O-alkyl-(N-imidazole), S-alkenyl, S-alkynyl, NH-alkyl, NH-alkenyl, NH-alkynyl, N-dialkyl, O-aryl, S-aryl, NH-aryl, O-aralkyl, S-aralkyl, NH-aralkyl, N-phthalimido, halogen (particularly fluoro), keto, carboxyl, nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, imidazole, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocycle, carbocycle, polyamine, polyamide, polyalkylene glycol, and polyethers of the formula (O-alkyl)_m, where m is 1 to about 10. Preferred among these polyethers are linear and cyclic polyethylene glycols (PEGs), and (PEG)-containing groups, such as crown ethers and those which are disclosed by Ouchi et al. (*Drug Design and Discovery* **1992**, 9, 93), Ravasio et al. (*J. Org. Chem.* **1991**, 56, 4329) and Delgado et. al. (*Critical Reviews in Therapeutic Drug Carrier Systems* **1992**, 9, 249), each of which is herein incorporated by reference in its entirety. Further sugar modifications are disclosed in Cook, P.D., *Anti-Cancer Drug Design*, **1991**, 6, 585-607. Fluoro, O-alkyl, O-alkylamino, O-alkyl imidazole, O-alkylaminoalkyl, and alkyl amino substitution is described in United States Patent Application serial number 08/398,901, filed March 6, 1995, entitled Oligomeric Compounds having Pyrimidine Nucleotide(s) with 2' and 5' Substitutions, hereby incorporated by reference in its entirety.


Additional substituent groups amenable to the present invention include -SR and -NR₂ groups, wherein each R is, independently, hydrogen, a protecting group or substituted or unsubstituted alkyl, alkenyl, or alkynyl. 2'-SR nucleosides are disclosed in United States Patent No. 5,670,633, issued September 23, 1997, hereby incorporated by reference in its entirety. The incorporation of 2'-SR monomer synthons are disclosed by Hamm et al., *J. Org. Chem.*, **1997**, 62, 3415-3420. 2'-NR₂ nucleosides are disclosed by Goettingen, M., *J. Org. Chem.*, **1996**, 61, 6273-6281; and Polushin et al., *Tetrahedron Lett.*, **1996**, 37, 3227-3230.

Further substituent groups have one of formula I or II:



wherein:

- 15 Z₀ is O, S or NH;
 J is a single bond, O or C(=O);
 E is C₁-C₁₀ alkyl, N(R₁)(R₂), N(R₁)(R₅), N=C(R₁)(R₂),
 N=C(R₁)(R₅) or has one of formula III or IV;



- 20 each R₆, R₇, R₈, R₉ and R₁₀ is, independently, hydrogen, C(O)R₁₁, substituted or unsubstituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, alkylsulfonyl, arylsulfonyl, a chemical

functional group or a conjugate group, wherein the substituent groups are selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl;

5 or optionally, R_7 and R_8 , together form a phthalimido moiety with the nitrogen atom to which they are attached;

or optionally, R_9 and R_{10} , together form a phthalimido moiety with the nitrogen atom to which they are attached;

each R_{11} is, independently, substituted or unsubstituted
10 C_1 - C_{10} alkyl, trifluoromethyl, cyanoethyloxy, methoxy, ethoxy, t-butoxy, allyloxy, 9-fluorenylmethoxy, 2-(trimethylsilyl)-ethoxy, 2,2,2-trichloroethoxy, benzyloxy, butyryl, iso-butyl, phenyl or aryl;

R_5 is T-L,

15 T is a bond or a linking moiety;

L is a chemical functional group, a conjugate group or a solid support material;

each R_1 and R_2 is, independently, H, a nitrogen protecting group, substituted or unsubstituted C_1 - C_{10} alkyl,
20 substituted or unsubstituted C_2 - C_{10} alkenyl, substituted or unsubstituted C_2 - C_{10} alkynyl, wherein said substitution is OR_3 , SR_3 , NH_3^+ , $N(R_3)(R_4)$, guanidino or acyl where said acyl is an acid amide or an ester;

or R_1 and R_2 , together, are a nitrogen protecting group
25 or are joined in a ring structure that optionally includes an additional heteroatom selected from N and O;

or R_1 , T and L, together, are a chemical functional group;

each R_3 and R_4 is, independently, H, C_1 - C_{10} alkyl, a
30 nitrogen protecting group, or R_3 and R_4 , together, are a nitrogen protecting group;

or R_3 and R_4 are joined in a ring structure that optionally includes an additional heteroatom selected from N and O;

033433 620430

Z_4 is OX, SX, or $N(X)_2$;

each X is, independently, H, C_1 - C_8 alkyl, C_1 - C_8 haloalkyl, $C(=NH)N(H)R_5$, $C(=O)N(H)R_5$ or $OC(=O)N(H)R_5$;

R_5 is H or C_1 - C_8 alkyl;

5 Z_1 , Z_2 and Z_3 comprise a ring system having from about 4 to about 7 carbon atoms or having from about 3 to about 6 carbon atoms and 1 or 2 hetero atoms wherein said hetero atoms are selected from oxygen, nitrogen and sulfur and wherein said ring system is aliphatic, unsaturated aliphatic,
10 aromatic, or saturated or unsaturated heterocyclic;

Z_5 is alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, aryl having 6 to about 14 carbon atoms, $N(R_1)(R_2)OR_1$, halo, SR_1 or CN;

15 each q_1 is, independently, an integer from 1 to 10;
each q_2 is, independently, 0 or 1;
 q_3 is 0 or an integer from 1 to 10;
 q_4 is an integer from 1 to 10;
 q_5 is from 0, 1 or 2; and

20 provided that when q_3 is 0, q_4 is greater than 1.

Representative substituent groups of Formula I are disclosed in United States Patent Application Serial No. 09/130,973, filed August 7, 1998, entitled "Capped 2'-Oxyethoxy Oligonucleotides," hereby incorporated by
25 reference in its entirety.

Representative cyclic substituent groups of Formula II are disclosed in United States Patent Application Serial No. 09/123,108, filed July 27, 1998, entitled "RNA Targeted 2'-Modified Oligonucleotides that are Conformationally
30 Preorganized," hereby incorporated by reference in its entirety.

Particularly preferred substituent groups include $O[(CH_2)_nO]_mCH_3$, $O(CH_2)_nOCH_3$, $O(CH_2)_nNH_2$, $O(CH_2)_nCH_3$, $O(CH_2)_nONH_2$, $O(CH_2)_nON[(CH_2)_nCH_3]_2$ (where n and m are from 1 to about 10),

C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino and substituted silyl. Another particularly preferred modification includes 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃ or 2'-MOE, Martin et al., *Helv. Chim. Acta*, 1995, 78, 486). A further preferred substituent group is 2'-dimethylamino-oxyethoxy, i.e., a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE. Representative aminoxy substituent groups are described in co-owned United States Patent Application serial number 09/344,260, filed June 25, 1999, entitled "Aminoxy-Functionalized Oligomers"; and United States Patent Application serial number 09/370,541, filed August 9, 1999, also identified by attorney docket number ISIS-3993, entitled Aminoxy-Functionalized Oligomers and Methods for Making Same; hereby incorporated by reference in their entirety.

Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on nucleosides and oligomers, particularly the 3' position of the sugar on the 3' terminal nucleoside or at a 3'-position of a nucleoside that has a linkage from the 2'-position such as a 2'-5' linked oligomer and at the 5'-position at a 5'-terminus. Oligomers may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugars structures include, but are not limited to, U.S. Patents 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned, and each of which is herein incorporated by

reference, and commonly owned United States patent application 08/468,037, filed on June 5, 1995, also herein incorporated by reference.

Representative guanidino substituent groups that are
5 shown in formula III and IV are disclosed in co-owned United States Patent Application 09/349,040, entitled "Functionalized Oligomers", filed July 7, 1999, hereby incorporated by reference in its entirety.

Representative acetamido substituent groups are
10 disclosed in United States Patent Application 09/378,568, entitled "2'-O-Acetamido Modified Monomers and Oligomers", filed August 19, 1999, also identified by attorney docket number ISIS-4071, hereby incorporated by reference in its entirety.

Representative dimethylaminoethoxyethyl substituent
15 groups are disclosed in International Patent Application PCT/US99/17895, entitled "2'-O-Dimethylaminoethoxyethyl-Modified Oligonucleotides", filed August 6, 1999, also identified by attorney docket number ISIS-4045, hereby
20 incorporated by reference in its entirety.

The use of mixed modifications in the terminal regions of an oligonucleotide to impart nuclease resistance is also within the scope of the present invention. For example an oligomeric compound of the invention can have enhanced
25 nuclease resistance resulting from one or more modified internucleoside linkages at the 5' end and one or more substituent groups at the 3' end. Another type of a mixed modification includes having a modified internucleoside linkage and a substituent group at the same end of a selected
30 oligomeric compound. Other examples include substituent groups or modified linkages used in conjunction with a non-standard linkage such as a 2', 5'-internucleoside linkage.

Oligomeric compounds according to the present invention preferably comprise from about 5 to about 50 nucleosides. It

0349739 " 64204960

is more preferred that such compounds comprise from about 8 to about 30 nucleosides, with 15 to 25 nucleosides being particularly preferred.

In general, the term "hetero" denotes an atom other than carbon, preferably but not exclusively N, O, or S. Accordingly, the term "heterocyclic ring" denotes an alkyl ring system having one or more heteroatoms (i.e., non-carbon atoms). Heterocyclic ring structures of the present invention can be fully saturated, partially saturated, unsaturated or with a polycyclic heterocyclic ring each of the rings may be in any of the available states of saturation. Heterocyclic ring structures of the present invention also include heteroaryl, which includes fused systems including systems where one or more of the fused rings contain no heteroatoms. Heterocycles, including nitrogen heterocycles, according to the present invention include, but are not limited to, imidazole, pyrrole, pyrazole, indole, 1H-indazole, α -carboline, carbazole, phenothiazine, phenoxazine, tetrazole, triazole, pyrrolidine, piperidine, piperazine and morpholine groups. A more preferred group of nitrogen heterocycles includes imidazole, pyrrole, indole, and carbazole groups.

A heterocyclic base moiety (often referred to in the art simply as a "base" or a "nucleobase") amenable to the present invention includes both naturally and non-naturally occurring nucleobases. The heterocyclic base moiety further may be protected wherein one or more functionalities of the base bears a protecting group. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine and guanine, and the pyrimidine bases thymine, cytosine and uracil. Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine

and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in the *Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., *Angewandte Chemie, International Edition*, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993.

Certain nucleobases are particularly useful for increasing the binding affinity of oligomeric compounds. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2 °C (*Id.*, pages 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-methoxyethyl sugar modifications.

Representative United States patents that teach the preparation of modified nucleobases include, but are not limited to, U.S. Patents 3,687,808; 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540;

5,587,469; 5,594,121, 5,596,091; 5,614,617; and 5,681,941, certain of which are commonly owned, and each of which is herein incorporated by reference, and commonly owned United States patent application 08/762,488, filed on December 10, 5 1996, also herein incorporated by reference.

The attachment of conjugate groups to oligomers is well documented in the prior art. The present methods include preparation of oligomeric compounds that include conjugate groups covalently bound to functional groups such as primary
10 or secondary hydroxyl groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic
15 properties of oligomers. Typical conjugates groups include cholesterols, phospholipids, biotin, phenazine, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention,
20 include groups that improve oligomer uptake, enhance oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution,
25 metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992, United States Patent No. 5,578,718, issued July 1, 1997, and United States Patent No. 5,218,105. Each of the foregoing is commonly assigned with this
30 application. The entire disclosure of each is incorporated herein by reference.

Preferred conjugate groups amenable to the present invention include lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86,

009483" 6204960

6553), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765), a
5 thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 111; Kabanov et al., *FEBS Lett.*, 1990, 259, 327; Svinarchuk et al., *Biochimie*, 1993, 75, 49), a phospholipid, e.g., di-
10 hexadecyl-rac-glycerol or triethylammonium-1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651; Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14,
15 969), adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651), a palmityl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264, 229), or an octadecylamine or hexylamino-carbonyl-oxysterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923).
20 Other groups that can be attached to oligomeric compounds to modify antisense properties include RNA cleaving complexes, pyrenes, metal chelators, porphyrins, alkylators, hybrid intercalator/ligands and photo-crosslinking agents. RNA cleavers include o-phenanthroline/Cu complexes and
25 Ru(bipyridine)₃²⁺ complexes. The Ru(bpy)₃²⁺ complexes are believed to interact with nucleic acids and cleave nucleic acids photochemically. Metal chelators include EDTA, DTPA, and o-phenanthroline. Alkylators include compounds such as iodoacetamide. Porphyrins include porphine, its substituted
30 forms, and metal complexes. Pyrenes include pyrene and other pyrene-based carboxylic acids that could be conjugated using the similar protocols.


039709"64649550

As used herein, "polyamine" refers to a moiety containing a plurality of amine or substituted amine functionalities. Polyamines according to the present invention have at least two amine functionalities.

5 "Polypeptide" refers to a polymer comprising a plurality of amino acids linked by peptide linkages, and includes dipeptides and tripeptides. The amino acids may be naturally-occurring or non-naturally-occurring amino acids. Polypeptides according to the present invention comprise at
10 least two amino acids.

The methods of the present invention can employ activated phosphorus compositions in coupling reactions. As used herein, the term activated phosphorus composition includes activated phosphorus containing monomers or
15 oligomers that are reactive with a hydroxyl group of another monomeric or oligomeric compound to form a phosphorus-containing internucleotide linkage. Such activated phosphorus groups contain activated phosphorus atoms in P^{III} valence state. Such activated phosphorus atoms are known in
20 the art and include, but are not limited to, phosphoramidite and chiral auxiliary moieties. A preferred synthesis utilizes phosphoramidites as activated phosphorus groups. Additional activated phosphites are disclosed in Tetrahedron Report Number 309 (Beaucage and Iyer, *Tetrahedron*, 1992, 48,
25 2223-2311).

Representative activated phosphorus containing monomers or oligomers include those having the formula:



wherein

each Bx is, independently, a heterocyclic base moiety or a blocked heterocyclic base moiety; and

each R₁ is, independently, H, a blocked hydroxyl group, a sugar substituent group, or a blocked substituent group;

5 T₃ is an hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleoside or an oligonucleotide;

R₄ is N(L₁)L₂;

each L₁ and L₂ is, independently, C₁₋₆ straight or branched alkyl, or a C₅₋₇ cyclic aliphatic ring system;

10 or L₁ and L₂ are joined together to form a 4- to 13-membered heterocyclic ring system including the nitrogen atom to which L₁ and L₂ are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S; and

15 R₅ is X₁;

X₁ is Pg-O-, Pg-S-, C₁-C₁₀ straight or branched chain alkyl, CH₃(CH₂)_{nn}-O-, R₂R₃N- or a group remaining from coupling a chiral auxiliary;

nn is from 0 to 10;

20 Pg is CH₃, -CH₂CH₂CN, -C(CH₃)(CH₃)-CCl₃, -CH₂-CCl₃, -CH₂CH=CH₂, CH₂CH₂SiCH₃, 2-yl-ethyl phenylsulfonate, δ-cyanobutenyl, cyano p-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-trifluoroacetyl ethyl, acetoxy phenoxy ethyl, or a blocking
25 group;

each R₂ and R₃ is, independently, hydrogen, C₁-C₁₀ alkyl, cycloalkyl or aryl;

or optionally, R₂ and R₃, together with the nitrogen atom to which they are attached form a cyclic moiety that may
30 include an additional heteroatom selected from O, S and N; or

R₄ and R₅ together with the phosphorus atom to which R₄ and R₅ are attached form a chiral auxiliary.


Groups that are attached to the phosphorus atom of internucleotide linkages before and after oxidation (R₄ and

009780" 64204960


R₅) can include nitrogen containing cyclic moieties such as morpholine. Such oxidized internucleoside linkages include a phosphoromorpholidothioate linkage (Wilk et al., *Nucleosides and nucleotides*, 1991, 10, 319-322). Further cyclic moieties
5 amenable to the present invention include mono-, bi- or tricyclic ring moieties which may be substituted with groups such as oxo, acyl, alkoxy, alkoxycarbonyl, alkyl, alkenyl, alkynyl, amino, amido, azido, aryl, heteroaryl, carboxylic acid, cyano, guanidino, halo, haloalkyl, haloalkoxy,
10 hydrazino, ODMT, alkylsulfonyl, nitro, sulfide, sulfone, sulfonamide, thiol and thioalkoxy. A preferred bicyclic ring structure that includes nitrogen is phthalimido.

439733 "64604960

Some representative examples of R_4 and R_5 groups which are known to the art skilled and are amenable to the present invention are shown below:



5 further examples include:



Functional groups including substituent groups discussed above which may be located on heterocyclic base and sugar moieties are routinely blocked with protecting (blocking groups) during synthesis and subsequently deblocked. In 5 general, a blocking group renders a chemical functionality of a molecule inert to specific reaction conditions and can later be removed from such functionality in a molecule without substantially damaging the remainder of the molecule. See, Green and Wuts, *Protective Groups in Organic Synthesis*, 10 2d edition, John Wiley & Sons, New York, 1991. For example, amino groups can be blocked with nitrogen protecting groups such as phthalimido, 9-fluorenylmethoxycarbonyl (Fmoc), triphenylmethylsulfenyl, t-BOC or benzyl groups. Carboxyl groups can be protected as acetyl groups. Representative 15 hydroxyl protecting groups are described by Beaucage et al., *Tetrahedron* 1992, 48, 2223. Preferred hydroxyl protecting groups are acid-labile groups, such as the trityl, monomethoxytrityl, dimethoxytrityl, trimethoxytrityl, 9-phenylxanthin-9-yl (Pixyl) and 9-(p-methoxyphenyl)xanthin-9- 20 yl (MOX). Chemical functional groups can also be "blocked" by including them in a precursor form. Thus an azido group can be considered a "blocked" form of an amine as the azido group is easily converted to the amine. Further representative protecting groups utilized in oligonucleotide 25 synthesis are discussed in Agrawal et al., *Protocols for Oligonucleotide Conjugates*, Eds., Humana Press, New Jersey, 1994, Vol. 26, pp. 1-72.

Standard oligonucleotide synthesis using phosphite (P^{III}) chemistry involves treatment of the growing oligomer with a 30 deprotecting reagent to create a free hydroxyl position that is available for a further coupling reaction. Hydroxyl protecting groups are preferably removed using a weak acid. Dependant on the choice of protecting group the deprotecting reagent can be acidic, basic, neutral or fluoride mediated.

009760" 6204960

A representative list of deprotecting reagents amenable to the present methods includes without limitation protic acids used for removing acid labile protecting groups such as dichloro- and trichloroacetic acids, Lewis acids such as BF_3 -etherate, zinc bromide, AlCl_3 , TiCl_4 , $(\text{Et})\text{AlCl}$, $(i\text{-Bu})_2\text{AlCl}$ and other reagents such as ceric ammonium nitrate, 1,1,1,3,3,3-hexafluoro-2-propanol, and diethyloxomalonate. A preferred deprotecting reagent that is used routinely for example for the removal of various trityl protecting groups is 2-5% dichloroacetic acid in either dichloromethane or dichloroethane.

The use of blocking groups is common practice to protect or block reactive or functional groups that are typically located on or linked to nucleobases, internucleotide linkages and sugars. Generally, blocking groups are removed using conditions that are stronger than those encountered during the iterative elongation steps in oligomer synthesis. This allows for deprotection of hydroxyl groups and coupling without effecting blocked positions. As used herein, the term "blocking group" describes a group that is stable to the conditions that are used to deprotect groups such as hydroxyl groups during the iterative elongation steps of oligomeric compound synthesis. Blocking groups are generally removed after the desired length has been synthesized. Standard phosphoramidite chemistry frequently uses acid labile protecting groups on hydroxyls that are used for coupling steps and strong base labile blocking groups to block other reactive positions not used in coupling steps. Many examples of protecting and blocking groups are collectively described in for example Green and Wuts *ibid*. Preferred blocking groups are removed by treatment with base. Some representative base labile protecting groups include without limitation, Fmoc (E. Atherton and R.C. Sheppard in *The Peptides*, S. Udenfriend, J. Meienhofer, Eds., Academic

Press, Orlando, 1987, volume 9, p.1), and various substituted sulfonylethyl carbamates exemplified by the Nsc group (Samukov et al., *Tetrahedron Lett*, 1994, 35:7821; Verhart and Tesser, *Rec. Trav. Chim. Pays-Bas*, 1987, 107:621).

5 After synthesis the resulting oligomeric compound generally is cleaved from the solid support to obtain the free oligomer. The step of deprotecting the blocked 5'-O-hydroxyl is usually performed separately as this is generally accomplished using an acidic deblocking reagent. This step
10 is routinely performed after deprotection, cleavage and purification has been performed to enhance the purification process by keeping the terminal hydroxyl blocked. The deprotection and cleavage steps can be separated into separate steps or combined into a single step depending on
15 the particular protecting groups, solid support linking groups and the choice of reagent or reagents used. In a preferred embodiment the simultaneous deprotection and cleavage of the final oligomeric compound following synthesis is accomplished in one step using a solution of ammonium
20 hydroxide (NH_4OH (30%) for 15 hours at 60 °C, filtered, rinsed with ethanol/water (1/1, v/v), the combined solutions are evaporated to dryness under vacuum).

The purification of oligomeric compounds is generally by reversed phase high performance liquid chromatography (RP-
25 HPLC) performed on a Waters Nova-Pak C18 column (3.9x300 mm) using a Waters HPLC system (600E System Controller, 996 Photodiode Array Detector, 717 Autosampler). For analysis an acetonitrile (A)/0.1M triethylammonium acetate gradient is used: 5% to 35% A from 0 to 10 min, then 35% to 40% A from 10
30 to 20 min, then 40% to 95% A from 20 to 25 min, flow rate = 10 mL/min/50% A from 8 to 9 min, 9 to 26 min at 50%, flow rate = 1.0 mL/min, $t_R(\text{DMT-off})$ 10-11 min, $t_R(\text{DMT-on})$ 14-16 min. The DMT-on fractions are collected and are evaporated

in vacuum, redissolved in water and the DMT group removed as described below.

Removal of the final hydroxyl protecting group from the 5'-hydroxyl group is generally performed by treatment with an acidic solution such as acetic acid. The oligomeric compound is treated with the acidic solution for about 30 minutes at room temperature. The mixture is further treated with sodium acetate and cold ethanol followed by vortexing and cooling with dry ice. The precipitate is centrifuged, separated, washed and dried to give the final deprotected product.

The term "nucleoside" as used in connection with this invention refers to a monomeric unit made up of a heterocyclic base moiety joined to a sugar moiety or sugar mimetic through a glycosyl linkage. The term "nucleotide" refers to a nucleoside having a phosphate group on its 3' or 5' sugar hydroxyl group.

In the context of this invention, the terms "oligomer" and "oligomeric compound" refer to a plurality of naturally-occurring or non-naturally-occurring nucleosides joined together in a specific sequence. The terms "oligomer" and "oligomeric compound" include oligonucleotides, oligonucleotide analogs, oligonucleosides and chimeric oligomeric compounds where there are more than one type of internucleoside linkages dividing the oligomeric compound into regions. Whereas the term "oligonucleotide" has a well defined meaning in the art, the term "oligomeric compound" or "oligomer" is intended to be broader, inclusive of oligomers having all manner of modifications known in the art. Gapped or chimeric compounds are disclosed in for example, U.S. Patent No. 5,623,065, issued April 22, 1997, the contents of which are incorporated herein by reference.

As used herein, the term "oligonucleoside" includes oligomers or polymers containing two or more nucleoside subunits having a non-phosphodiester linking moiety.


Oligonucleosides according to the invention have a ribofuranose moiety attached to a nucleobase through a glycosyl bond.

Gapmer technology has been developed to incorporate
5 modifications at the ends ("wings") of oligomeric compounds, leaving a phosphorothioate gap in the middle for RNase H activation (Cook, P.D., *Anti-Cancer Drug Des.*, **1991**, 6, 585-607; Monia et al., *J. Biol. Chem.*, **1993**, 268, 14514-14522). In a recent report, the activities of a series of uniformly
10 2'-O modified 20 mer RNase H-independent oligonucleotides that were antisense to the 5'-cap region of human ICAM-1 transcript in HUVEC cells, were compared to the parent 2'-deoxy phosphorothioate oligonucleotide (Baker et al., *J. Bio. Chem.*, **1997**, 272, 11994-12000). The 2'-MOE/P=O oligomer
15 demonstrated the greatest activity with an IC₅₀ of 2.1 nM (T_m = 87.1°C), while the parent P=S oligonucleotide analog had an IC₅₀ of 6.5 nM (T_m = 79.2°C). Correlation of activity with binding affinity is not always observed as the 2'-F/P=S (T_m = 87.9°C) was less active than the 2'-MOE/P=S (T_m = 79.2°C) by
20 four fold. The RNase H competent 2'-deoxy P=S parent oligonucleotide exhibited an IC₅₀ = 41 nM.

As used herein the term "chiral auxiliary" is meant to include groups that function to provide chirality to internucleoside phosphorus linkages during synthesis. Chiral
25 auxiliaries amenable to the present invention include those that form a P^{III} intermediate capable of being oxidized. Chiral auxiliaries will give either Sp or Rp chirality for the respective internucleoside linkage in the final oligomeric compound. Accordingly, chiral auxiliaries are
30 allowed to remain on the growing chain, and are removed at the end of the iterative synthetic regime. Removal of chiral auxiliaries can be conveniently accomplished in a single treatment after the completion of the iterative synthesis. Chiral auxiliaries and methods of their incorporation using

standard protocols are disclosed in commonly owned United States patent application 09/438,989, filed on November 12, 1999, incorporated herein by reference. Further chiral auxiliaries have been previously reported for use in the
5 preparation of oligomeric phosphorothioates (see Iyer et al., *Tetrahedron letters*, **1998**, 39, 2491-2494 and Wilk et al., *J. Am. Chem. Soc.*, **2000**, 122, 2149-2156). Representative chiral auxiliaries include, without limitation those having the following formulas:

10



As used herein, the term "alkyl" includes, but is not limited to, straight chain, branched chain and alicyclic hydrocarbon groups. Alkyl groups of the present invention may be substituted. Representative alkyl substituents are
15 disclosed in United States Patent No. 5,212,295, at column 12, lines 41-50, hereby incorporated by reference in its entirety. Substituent groups include, but are not limited to, alkyl, alkenyl, alkynyl, aryl, hydroxyl, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, thioalkyl,
20 trifluoromethyl, halo, nitrile, trifluoromethoxy and azido. As used herein, the term "lower alkyl" is intended to mean an alkyl group having 10 or fewer carbons.

Alkenyl groups according to the invention are to straight chain, branch chain, and cyclic hydrocarbon groups
25 containing at least one carbon-carbon double bond, and

alkynyl groups are to straight chain, branch chain, and cyclic hydrocarbon groups containing at least one carbon-carbon triply bond. Alkenyl and alkynyl groups of the present invention can be substituted.

5 Aryl groups are substituted and unsubstituted aromatic cyclic moieties including but not limited to phenyl, naphthyl, anthracyl, phenanthryl, pyrenyl, and xylyl groups. Alkaryl groups are those in which an aryl moiety links an alkyl moiety to a core structure, and aralkyl groups are
10 those in which an alkyl moiety links an aryl moiety to a core structure.

As used herein, the term "aralkyl" denotes alkyl groups which bear aryl groups, for example, benzyl groups. The term "alkaryl" denotes aryl groups which bear alkyl groups, for
15 example, methylphenyl groups. As used herein, the term "aryl" denotes aromatic cyclic groups including, but not limited to, phenyl, naphthyl, anthracyl, phenanthryl and pyrenyl. Preferred aryl and aralkyl groups include, but are not limited to, phenyl, benzyl, xylyl, naphthyl, toluy,
20 pyrenyl, anthracyl, azulyl, phenethyl, cinnamyl, benzhydryl, and mesityl. Typical substituents for substitution include, but are not limited to, hydroxyl, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, or alkyl, aryl, alkenyl, or alkynyl groups.

25 As used herein, the term "alkanoyl" has its accustomed meaning as a group of formula $-C(=O)-alkyl$. A preferred alkanoyl group is the acetyl group.

In one aspect of the present invention oligomeric compounds are prepared using support bound methodologies
30 wherein the oxidation and capping steps are combined into a single step. A first modified or unmodified nucleoside is attached to a support media preferably via a linkage to the 3'-position. The nucleoside could alternatively be attached to a support media through the 2'-position as when preparing

039480" 02204960

positionally modified internucleoside linkages. Alternatively, the support media with the desired nucleoside attached can be purchased from a number of commercial sources. In a traditional synthesis this nucleoside will ultimately become
5 the nucleoside at the 3'-end of the final oligomeric compound. The support media with the nucleoside attached is placed in a reaction vessel such as a glass reactor. One of the hydroxyl groups (preferably the 5'-hydroxyl group) is deprotected and treated with a second nucleoside having a
10 group reactive with the hydroxyl group such as an activated phosphorus group or a chiral auxiliary. This coupling step is preferably performed in the presence of an activating agent such as DBU or 1-H-tetrazole. The linkage thus formed is treated with a mixture containing reagents for oxidizing
15 and capping. A preferred mixture for incorporating a sulfur atom is (0.3M) dimethylthiuram disulfide in cap A (20% acetic anhydride in acetonitrile) mixed with and equal volume of cap B (20% N-methylimidazole, 30% pyridine and 50% acetonitrile, by volume). The cycle is optionally repeated to add
20 additional nucleosides until the desired oligomeric compound is completed.

As used herein, the term "sulfurizing reagent" includes without limitation, dimethylthiuram disulfide (Cummings et al., *Ind. Eng. Chem.*, **1928**, 20, 1173); 1,2,4-dithiazolidine-
25 3,5-dione (DTSNH, see Xu et al., *Nucleic Acids Research*, **1996**, 24, 1602-1607); 3-methyl-1,2,4-dithiazolin-5-one (MEDITH, see Zang et al., *Tetrahedron Lett.*, **1999**, 40, 2095-20980); phenylacetyl disulfide (PADS, see Kamer et al., *Tetrahedron Lett.*, **1989**, 30, 6757-6760; Cheruvallath et al.,
30 *Organic Process Research & Development*, **2000**, 4, 199-204); tetraethylthiuram disulfide (TETD, see Vu et al., *Tetrahedron Lett.*, **1991**, 32, 3005-3008); 3H-1,2-benzodithiol-3-one-1,1-dioxide (Beaucage reagent, see e.g. Iyer, R.P., et.al., J.

Chem. Soc., 1990, 112, 1253-1254, and Iyer, R.P., et.al., J. Org. Chem., 1990, 55, 4693-4699); tetraethylthiuram disulfide (see e.g., Vu, H., Hirschbein, B.L., *Tetrahedron Lett.*, 1991, 32, 3005-3008); dibenzoyl tetrasulfide (see e.g., Rao, M.V., et.al., *Tetrahedron Lett.*, 1992, 33, 4839-4842); benzyltriethylammonium tetrathiomolybdate (BTM, see e.g., Rao, M.V., et.al., *Tetrahedron Lett.*, 1994, 35, 6741-6744); di(phenylacetyl)disulfide (see e.g., Kamer, P.C.J., *Tetrahedron Lett.*, 1989, 30, 6757-6760); Bis(O,O-diisopropoxy phosphinothioyl)disulfides (see Stec et al., *Tetrahedron Lett.*, 1993, 34, 5317-5320); 3-ethoxy-1,2,4-dithiazoline-5-one (EDITH, see Xu et al., *Nucleic Acids Research*, 1996 24, 1602-1607, and *Nucleic Acids Research*, 1996 24, 3643-3644); Bis(p-chlorobenzenesulfonyl)disulfide (see *Nucleic Acids Research*, 1995 23, 4029-4033); bis(ethoxythiocarbonyl)-tetrasulfide (see Zang et al., *Tetrahedron Lett.*, 1998, 39, 2467-2470); bis(p-toluenesulfonyl)disulfide (Efimov et al., *Nucleic Acids Res.*, 1995, 23, 4029-4033); 3-amino-1,2,4-dithiazole-5-thione (see *Org. Process Res. Dev.*, 2000, 4, 194-198); ethylthiuram disulfide CAS # 3082-38-0; 5,6-dihydro-3H-imidazo[2,1-C]-1,2,4-dithiazole-3-thione CAS # 33813-20-6; 4-methyl-5-(methylimino)-1,2,4-dithiazolidine-3-thione CAS # 20042-85-7; sulfur, sulfur in combination with ligands like triaryl, trialkyl, triaralkyl, or trialkaryl phosphines. The foregoing references are hereby incorporated by reference in their entirety.

A preferred list of sulfurizing reagents includes: 3-amino-1,2,4-dithiazole-5-thione; 3-ethoxy-1,2,4-dithiazoline-5-one; 1,2,4-dithiazolidine-3,5-dione; 3-methyl-1,2,4-dithiazolin-5-one; and dimethylthiuram disulfide.

A representative list of capping reagents useful in the methods of the present invention include without limitation, acetic anhydride, *t*-butylphenoxyacetic anhydride, phosphite

monoesters, and selected acid chlorides preferably delivered concurrently with a nucleophilic catalyst (e.g. a strong base) such as for example N-methylimidazole or triethylamine. Generally capping reagents comprise a mixture of Cap A and

5 Cap B. Representative mixtures include without limitation:

Cap A: acetic anhydride in acetonitrile or tetrahydrofuran;

chloroacetic anhydride in acetonitrile or tetrahydrofuran;

10 Cap B: N-methylimidazole and pyridine in acetonitrile or tetrahydrofuran;

4-dimethylaminopyridine (DMAP) and pyridine in acetonitrile or tetrahydrofuran;

2,6-lutidine and N-methylimidazole in

15 acetonitrile or tetrahydrofuran.

A more detailed description capping reagents is discussed in United States Patent 4,816,571, issued March 28, 1989 which is incorporated herein by reference. A preferred capping reagent is acetic anhydride routinely used as a

20 mixture of cap A and cap B.

During the coupling step one compound having an active phosphate is coupled to a second compound having a free hydroxyl group. An activating agent is not believed to be essential for this step but one is generally used to increase

25 the reaction efficiency. A list of activators and references for each can be found in Eleueri et al., *Organic Process Research & Development*, 2000, 4, 182-189. Preferred activators include without limitation: 1H-tetrazole, 5-(2-nitrophenyl)-1H-tetrazole, 5-(p-nitrophenyl)-1H-tetrazole, 5-

30 trifluoromethyl-1H-tetrazole, 5-ethylthio-1H-tetrazole, 5-benzylthio-1H-tetrazole, 2,4,5-tribromoimidazole, 2-nitroimidazole, 4,5-dichloroimidazole, 2-bromo-4,5-dicyanoimidazole, 4,5-dicyanoimidazole, N-methylimidazole hydrochloride, 1-hydroxybenzotriazole, 5-chlorobenzotriazole,

chlorotrimethylsilane, benzimidazolium triflate, imidazolium triflate, pyridinium hydrochloride/imidazole, pyridinium tetrafluoroborate, pyridinium chloride, pyridinium bromide, pyridinium 4-methylbenzinesulfonate, 5 N-methylimidazolium trifluoroborate, N-methylanilinium trichloroacetate, N-methylanilinium trifluoroacetate, 1H-tetrazole/DMAP, 1H-tetrazole/N-methylimidazole, and N-methylimidazolium trifluoromethanesulfonate (see review article Beaucage et al., *Current Protocols in Nucleic Acid* 10 *Chemistry*, **2000**, 3.3.1-3.3.20).

The current method of choice for the preparation of oligomeric compounds uses support media. Support media is used to attach a first nucleoside or larger nucleosidic synthon which is then iteratively elongated to give a final 15 oligomeric compound. Support media can be selected to be insoluble or have variable solubility in different solvents to allow the growing oligomer to be kept out of or in solution as desired. Traditional solid supports are insoluble and are routinely placed in a reaction vessel while 20 reagents and solvents react and or wash the growing chain until cleavage frees the final oligomer. More recent approaches have introduced soluble supports including soluble polymer supports to allow precipitating and dissolving the bound oligomer at desired points in the synthesis (Gravert et 25 al., *Chem. Rev.*, **1997**, 97, 489-510). Representative support media that are amenable to the methods of the present invention include without limitation: controlled pore glass (CPG); oxalyl-controlled pore glass (see, e.g., Alul, et al., *Nucleic Acids Research* **1991**, 19, 1527); TENTAGEL Support, 30 (see, e.g., Wright, et al., *Tetrahedron Letters* **1993**, 34, 3373); or POROS, a copolymer of polystyrene/divinylbenzene available from Perceptive Biosystems. The use of a soluble support media, poly(ethylene glycol), with molecular weights between 5 and 20 kDa, for large-scale synthesis of

phosphorothioate oligonucleotides is described in, Bonora et al., *Organic Process Research & Development*, 2000, 4, 225-231.

It was previously reported (Cummings A. D. et al. 5 Ind. Eng. Chem., 1928, 20, 1173) that dimethylthiuram disulfide was not a stable compound and decomposes slowly on standing. The dimethylthiuram disulfide, after standing for 1 month, failed to show a melting point of 102°C. The decomposition products were identified as hydrogen sulfide, 10 elemental sulfur and methyl isothiocyanate. The lack of long shelf life for dimethylthiuram disulfide has been attributed to the dithiocabamate derivative of methylamine, which is a primary amine. The present invention provides a new and improved procedure for the synthesis of dimethylthiuram 15 disulfide (see Example 7) which utilizes an acid wash at the end of the synthesis. The primary amine is protonated which stabilizes the dithiocarbamate structure from degradation. A further improvement was realized by oxidizing the intermediate with hydrogen peroxide as opposed to iodine. This 20 led to the preparation of white crystalline product instead of the yellow unstable product previously reported. Dimethylthiuram disulfide made by this protocol is very stable even after six months of storage at room temperature.

Oligomeric compounds prepared by the methods of the 25 present invention can be used in diagnostics, therapeutics and as research reagents and kits. They can also be used in pharmaceutical compositions by including a suitable pharmaceutically acceptable diluent or carrier. They can further be used for treating organisms having a disease characterized 30 by the undesired production of a protein. For this purpose, the organism is contacted with an oligomer having a sequence that is capable of specifically hybridizing with a strand of nucleic acid encoding the undesirable protein. Treatments of this type can be practiced on a variety of organisms ranging

009763 02204960

from unicellular prokaryotic and eukaryotic organisms to multicellular eukaryotic organisms. Any organism that utilizes DNA-RNA transcription or RNA-protein translation as a fundamental part of its hereditary, metabolic or cellular control is susceptible to therapeutic and/or prophylactic treatment in accordance with the invention. Seemingly diverse organisms such as bacteria, yeast, protozoa, algae, all plants and all higher animal forms, including warm-blooded animals, can be treated. Further, each cell of multicellular eukaryotes can be treated, as they include both DNA-RNA transcription and RNA-protein translation as integral parts of their cellular activity. Furthermore, many of the organelles (e.g., mitochondria and chloroplasts) of eukaryotic cells also include transcription and translation mechanisms. Thus, single cells, cellular populations or organelles can also be included within the definition of organisms that can be treated with therapeutic or diagnostic oligomeric compounds of the invention.

Those skilled in the art will appreciate that numerous changes and modifications may be made to the preferred embodiments of the invention and that such changes and modifications may be made without departing from the spirit of the invention. It is, therefore, intended that the appended claims cover all such equivalent variations as fall within the true spirit and scope of the invention.

EXAMPLE 1

5c-Methyl-2t [(1-methyl-1-methylamino) ethyl]-cyclohexan-1r-ol (Compound 1)


The title compound is synthesized according to a literature procedure using (+)-pulegone (He et al., *J. Org. Chem.*, 1990, 55, 2114-2119) by first preparing 5c-Methyl-2t [(1-methyl-1-benzylamino) ethyl]-cyclohexan-1r-ol. This compound is subjected to hydrogenolysis by Pd/H₂ to give the

corresponding amino alcohol (removal of benzyl group). The amino alcohol is then treated with 1 equivalent of HCHO followed by NaCNBH₃ reduction to give the title Compound. This isomer is used to prepare Rp phosphorothioate linkages.

5 The isomer of the title compound (Compound 2) is prepared from the naturally occurring (-)-pulegone (available from Fluka), Compound 2 is obtained as a Chiral Adjuvant following a literature procedure (He et al., *Tetrahedron*, 1987, 43, 4979-4987). This isomer is used to prepare Sp
10 phosphorothioate linkages.

Example 2


Preparation of Sp Monomer (Compound 3)



Compound 2 is treated with PCl₃ (1 equivalent) with
15 excess of Hunig base in THF solvent at -5°C for 10 minutes. The resulting chloro compound is treated with a selected 2'-deoxy-5-O-DMT-nucleoside having a free 3'-OH group (2'-O-deoxy-5'-O-DMT-6-N-benzoyl adenosine, 2'-O-deoxy-5'-O-DMT-4-N-benzoyl cytosine, 2'-O-deoxy-5'-O-DMT-2-N-butyryl
20 guanosine, 2'-O-deoxy-5'-O-DMT-thymidine or modified optionally protected 5-O-DMT-nucleoside). TLC and ¹³C NMR analysis is used to reveal the formation of a single diastereomer. The crude material is washed with saturated sodium bicarbonate and dried over anhydrous sodium sulfate.
25 The resulting material is purified either by crystallization or by silica gel column chromatography.

EXAMPLE 3**protected dimer, (Compound 4)**

Purified compound 3 is condensed with a 5'-HO-T-CPG (Example 5), or other solid support bound 5'-OH-nucleoside (such as 2'-O-deoxy-6-N-benzoyl adenosine, 2'-O-deoxy-4-N-benzoyl cytidine, 2'-O-deoxy-2-N-isobutyryl guanosine or other modified optionally protected 5'-OH'-3'-CPG-nucleoside), for 2 hours using tetrazole as the coupling agent. The capping and sulfurization is completed in one step using (0.3M) dimethylthiuram disulfide in cap A (20% acetic anhydride in acetonitrile) mixed with an equal volume of cap B (20% N-methylimidazole, 30% pyridine and 50% acetonitrile, by volume) giving the protected phosphorothioate dimer attached to solid support. The protected dimer is cleaved from the solid support, deprotected by treatment with concentrated ammonium hydroxide (30%, 12 hours), and purified by HPLC. The nucleoside dimer is treated with 80% aqueous acetic acid to remove the 5'-triyl group. The Sp configuration is assigned as described below in the procedures.

EXAMPLE 4**Preparation of Rp Monomer (Compound 5)**

The Rp monomer is prepared following the procedures illustrated for the Sp dimer in example 2 using Compound 1.

EXAMPLE 5

Preparation of Rp Dimer (Compound 6)

The Rp dimer is prepared following the procedures illustrated for the Sp dimer in example 3 using Compound 5.

EXAMPLE 6

5 **Synthesis of Chirally pure 5'-T_{Sp}T_{Rp}T_{Rp}T_{Rp}T_{Rp}T_{Sp}T-3' phosphorothioate heptamer**

50 milligram (2 μ mole) of 5'-O-dimethoxytritylthymidine bound to CPG (controlled pore glass) through an ester linkage is taken up in a glass reactor, and a toluene solution of 3%
10 dichloroacetic acid (v/v) is added to deprotect the 5'-hydroxyl group. The product is washed with acetonitrile and a 0.2 M solution of Compound 3 (B=T) in acetonitrile (25 fold excess) and a 0.5 M solution of DBU in acetonitrile (200 fold excess) is added and allowed to react at room temperature for
15 15 minutes. The product is washed with acetonitrile followed by the addition of a solution of (0.3M) dimethylthiuram disulfide in cap A (20% acetic anhydride in acetonitrile) mixed with and equal volume of cap B (20% N-methylimidazole, 30% pyridine and 50% aetonitrile, by volume) with reaction
20 allowed to progress at room temperature for 5 minutes. The product is washed with acetonitrile.

In the next cycle Compound 5 (B=T) is used as the incoming monomer and the cycle is repeated. This complete cycle is repeated four more times to introduce the Rp
25 linkages. In the final cycle Compound 3 is used as the incoming monomer which introduces the terminal Sp linkage. The solid support containing the heptamer is treated with 30% aqueous ammonium hydroxide solution for 90 minutes at room temperature. The aqueous solution is filtered, and
30 concentrated under reduced pressure to give the chirally pure phosphorothioate heptamer.

EXAMPLE 7

Preparation of dimethylthiuram disulfide

In a 4-liter bottle equipped with a mechanical stirrer, sodium hydroxide (80 g, 2 mol) was dissolved in water (500 mL) and the solution was cooled to 0°C (ice-water bath). THF (200 mL), methylamine (40% in water, 170 mL, 2 mol) and carbon disulfide (120 mL, 2 mol) were added and the mixture was stirred at 0°C for 30 minutes. Crushed ice (1.5 kg) was added, followed by glacial acetic acid (300 mL). Hydrogen peroxide (30%, 100 mL, 1 mol) was added dropwise over 10 minutes with temperature maintained below 5°C by adding ice with stirring. Hexanes (or heptane) (800 mL) was added and the mixture was stirred for another 30 minutes in an ice-water bath. The mixture was filtered and washed with aqueous trichloroacetic acid (2%, 5x200 mL) and hexanes (or heptane) (2x200mL). The product was dried in air for 1 day to a constant weight to give an off-white solid (205.8, yield: 97%). M.p. 98-100°C (dec.)

HPLC analysis:

Column: YMC ODS-AQ S3 120A, 4.6x100mm
Flow rate: 1.0 mL/min
Detector: UV at 244 nm
Sample: inject 10 µL (1-2 mg in 1 mL of glacial acetic acid)
Retention time: 5.4 min

25


Linear gradient

<u>Time (Min)</u>	<u>Acetonitrile</u>	<u>0.2% Acetic acid</u>
0	50	50
15	90	10
25	90	10
30	50	50
35	50	50

The crude product (200g) was recrystallized by dissolution in THF (1L) containing trichloroacetic acid (10 g). Water (200 mL) was added followed by slow addition of hexanes (5 L) over 30 minutes. After stirring at 0°C for 15 hour, filtering, washing with 2% aqueous trichloroacetic acid (3x200 mL) followed by drying the recrystallized dimethylthiuram disulfide was obtained.

EXAMPLE 8**Sulfurization of triethyl phosphite with various sulfurizing****10 reagents**

Small scale sulfurization reactions were performed in NMR tubes using CD₃CN to measure the efficiency of various sulfurization reagents. The basic reaction scheme is shown below:



	Reagent (=S)	% P=O	% product	% product (w/tetrazole)
	Beaucage	0.7	100	---
20	Phenylacetyl disulfide	0.7	100	---
	Tetraethylthiuram disulfide	2	82	100
	Tetramethylthiuram disulfide	4	2	93
25	Morpholino thiocarbonyl disulfide	1.7	50	93
	Dimethylthiuram disulfide	0	100	100

30 The experimental conditions varied dependent on the sulfurizing reagent used. The conditions are detailed below:

1. 0.5 M beaucage in CD₃CN (1mL) and triethyl phosphite (17μL) for 5 min.

003483 04204960

2. 0.2 M PADS in 3-picoline-CD₃CN (1:1, 1mL) and triethyl phosphite (17μL) for 5 min.

3. 0.2 M tetraethylthiuram disulfide in CD₃CN (1mL) and triethyl phosphite (17μL) for 5 min.

5 4. 0.2 M tetraethylthiuram disulfide and 0.3 M tetrazole in CD₃CN (1mL) and triethyl phosphite (17 μL) for 5 min.

Example 9

Solid phase synthesis of full phosphorothioate 5'-TTTTTTC-3' using tetraethylthiuram disulfide

The above sequence was synthesized following standard phosphoramidite protocols. The 4-step procedure was performed on via automated synthesis using a 6.3 mL column on an OligoPilot II (Armstrong Pharmacia). The reagents and the amounts used are as follows:

1. Detritylation: 3% dichloroacetic acid in dichloromethane.

2. Coupling: 2 equivalents of phosphoramidite for 5 min.

20 3. Thiolation: 1 column volume (CV) 0.5 M tetraethylthiuram disulfide and 0.45 M tetrazole in acetonitrile for 10 min.

4. Capping: 0.5 CV of cap A (20% acetic anhydride in acetonitrile) and cap B (N-methylimidazole-pyridine-acetonitrile, 2:3:5, v/v/v) for 1 min.

25 After synthesis, the solid support was heated with concentrated aqueous ammonia solution (50 mL) at 58°C overnight and the filtered. The filtrate was concentrated under reduced pressure and dried. The ³¹P NMR study showed
30 3.7% P=O and 96.3% P=S.

Example 10

Solid phase synthesis of full phosphorothioate 5'-TTTTTTC-3' using morpholino thiocarbonyl disulfide

The above sequence was synthesized following standard phosphoramidite protocols. The 4-step procedure was
5 performed on via automated synthesis using a 6.3 mL column on an OligoPilot II (Amersham Pharmacia). The reagents and the amounts used are as follows:

1. Detritylation: 3% dichloroacetic acid in dichloromethane.
- 10 2. Coupling: 2 equivalents of phosphoramidite for 5 min.
3. Thiolation: 1 column volume (CV) 0.3M morpholino thiocarbonyl disulfide and 0.23M tetrazole in dichloromethane-acetonitrile (1:1 v/v) for 10 min.
- 15 4. Capping: 0.5 CV of cap A (20% acetic anhydride in acetonitrile) and cap B (N-methylimidazole-pyridine-acetonitrile, 2:3:5, v/v/v) for 1 min.

After synthesis, the solid support was heated with concentrated aqueous ammonia solution (50 mL) at 58°C
20 overnight and the filtered. The filtrate was concentrated under reduced pressure and dried. ³¹P NMR showed 3.3% P=O and 96.7% P=S.

037433 "4407" 620430

EXAMPLE 11

Solid phase synthesis of full phosphorothioate 20mer, SEQ ID NO. 1 (5'-GTGCTCATGG TGCACGGTCT-3'; all C's are 5-Me-C's)

A. Using phenylacetyl disulfide (PADS)

5 The above sequence (SEQ ID NO. 1) was synthesized following standard phosphoramidite protocols. The 4-step procedure was performed on via automated synthesis using a 6.3 mL column on an OligoPilot II (Armstrong Pharmacia). The reagents and the amounts used are as follows:

- 10 1. Detritylation: 3% dichloroacetic acid in dichloromethane.
2. Coupling: 2 equivalents of phosphoramidite for 5 min.
3. Thiolation: 1 column volume (CV) 0.2 M phenylacetyl
15 disulfide (PADS) in 3-picoline-acetonitrile (1:1 v/v) for 10 min.
4. Capping: 0.5 CV of cap A (20% acetic anhydride in acetonitrile) and cap B (N-methylimidazole-pyridine-acetonitrile, 2:3:5, v/v/v) for 1 min.

20 After synthesis, the solid support was heated with concentrated aqueous ammonia solution (50 mL) at 58°C overnight and the filtered. The filtrate was concentrated under reduced pressure and dried. Purity and ³¹P NMR data are shown below.

25 B. Using dimethylthiuram disulfide

The above sequence (SEQ ID NO. 1) was synthesized following standard phosphoramidite protocols. The 3-step procedure was performed on via automated synthesis using a 6.3 mL column on an OligoPilot II (Armstrong Pharmacia).

- 30 1. Detritylation: 3% dichloroacetic acid in dichloromethane.
2. Coupling: 2 equivalents of phosphoramidite for 5 min.

3. Thiolation-capping: 1 column volume (CV) 0.3 M dimethylthiuram disulfide (DMDS) in cap A (20% acetic anhydride in acetonitrile) and 1CV cap B (N-methylimidazole-pyridine-acetonitrile, 2:3:5, v/v/v) for 1 min.

5 After synthesis, the solid support was heated with concentrated aqueous ammonia solution (50 mL) at 58°C overnight and the filtered. The filtrate was concentrated under reduced pressure and dried. Purity and ³¹P NMR data are shown below.

10

<u>Rgt</u>	<u>Loading</u>	<u>Crude O.D.</u>	<u>O.D./mmol</u>	<u>%Trityl-on</u>	<u>%CGE</u>	<u>%P=O</u>
PADS	173	25625	148	74.7	78.1	0.40
DMDS	169	26265	149	76.2	81.5	0.22.

15 **EXAMPLE 12**

Preparation of thymidine 8-mer having phosphodiester internucleotide linkages using a single step combining oxidation and capping (5'-TTTTTTTT-3')

20 A thymidine 8-mer was prepared following the procedures illustrated above (see example 11). The synthesis was performed as per the 3-step procedure (combined oxidation and capping steps) using a 6.3 mL column on an OligoPilot II (Armstrong Pharmacia) and 1.89 g Primer Support T (93 μmol/g).

- 25 1. Detritylation: 3% dichloroacetic acid in toluene.
2. Coupling: 3 equivalents of phosphoramidite for 5 min.
3. Oxidation-capping: 1 CV of 0.1 M iodine in cap A (20% acetic anhydride in acetonitrile) and 1 CV of cap B (N-methylimidazole-pyridine-acetonitrile, 2:3:5, v/v/v) for 2 min.
- 30

After synthesis, the solid support was heated with concentrated aqueous ammonia solution (50 mL) at 58°C

overnight and the filtered. The filtrate was concentrated under reduced pressure prior to analysis.


Crude yield: 7564 O.D.; full length percentage: 90.1%;
³¹P NMR: -0.383, -0.276 and -0.215 ppm.

11439734 "E" 204350

WHAT IS CLAIMED IS:

1. A method of preparing an oligomeric compound having at least one moiety of formula:

5



wherein:

X_2 is O or S;

X_1 is Pg-O-, Pg-S-, C_1 - C_{10} straight or branched chain alkyl, $CH_3(CH_2)_{nn}$ -O-, R_2R_3N - or a group remaining from coupling
10 a chiral auxiliary;

nn is from 0 to 10;

Pg is CH_3 , $-CH_2CH_2CN$, $-C(CH_3)(CH_3)-CCl_3$, $-CH_2-CCl_3$, $-CH_2CH=CH_2$, $CH_2CH_2SiCH_3$, 2-yl-ethyl phenylsulfonate, δ -cyano-butenyl, cyano *p*-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-
15 trifluoroacetyl ethyl, acetoxy phenoxy ethyl, or a blocking group;

each R_2 and R_3 is, independently, hydrogen, C_1 - C_{10} alkyl, cycloalkyl or aryl;

20 or optionally, R_2 and R_3 , together with the nitrogen atom to which they are attached form a cyclic moiety;


each Bx is, independently, a heterocyclic base moiety;

and

each R_1 is, independently, H, a blocked hydroxyl group,
25 or a sugar substituent group;

comprising the steps of:

(a) providing a 5'-O-protected compound of the formula:




wherein:

5 T_1 is a hydroxyl protecting group; and

T_2 is a covalent attachment to a support media, a nucleoside bound to a support media, a nucleotide, an oligonucleoside or an oligonucleotide;

(b) treating said 5'-O-protected compound with a
10 deprotecting reagent for a time and under conditions effective to form a 5'-O-deprotected compound;

(c) coupling said 5'-O-deprotected compound with an activated phosphorus composition of the formula:



15 wherein:

T_3 is a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleoside or an oligonucleotide;

R_4 is $N(L_1)L_2$;

each L_1 and L_2 is, independently, C_{1-6} straight or
20 branched alkyl, or a C_{5-7} cyclic aliphatic ring system;


or L_1 and L_2 are joined together to form a 4- to 13-membered heterocyclic ring system including the nitrogen atom to which L_1 and L_2 are attached; and

R_5 is X_1 ;

or R₄ and R₅ together with the phosphorus atom to which R₄ and R₅ are attached form a chiral auxiliary;

for a time and under conditions effective to form an extended compound having the formula:

5



(d) treating said extended compound with a mixture comprising an oxidizing reagent and a capping reagent for a time and under conditions effective to form said oligomeric compound.

10 2. The method of claim 1 further comprising treating said oligomeric compound with a reagent for a time and under conditions effective to remove said blocking groups thereby forming a deblocked oligomeric compound.

15 3. The method of claim 2 wherein said reagent is effective to cleave the oligomeric compound from the support media.

4. The method of claim 3 wherein said reagent is aqueous ammonium hydroxide.

20 5. The method of claim 2 further comprising treating said oligomeric compound with a further reagent for a time

"039430" 24204960

and under conditions effective to cleave the oligomeric compound from the support media.

6. The method of claim 1 further comprising treating said oligomeric compound with a deprotecting reagent for a time and under conditions effective to deprotect the T₃ hydroxyl protecting group.

7. The method of claim 1 wherein said mixture comprises from 0.02M to 0.2M oxidizing reagent.

8. The method of claim 7 wherein said mixture comprises from 0.1M to 0.2M oxidizing reagent.

9. The method of claim 1 wherein said oxidizing reagent transfers an oxygen atom.

10. The method of claim 9 wherein said oxidizing reagent is iodine, *m*-chloroperbenzoic acid, iodobenzene diacetate, tetra-*n*-butylammonium periodate, *tert*-butyl hydroperoxide, di-*tert*-butyl hydroperoxide, cumene hydroperoxide, hydrogen peroxide; bis-trimethylsilyl peroxide; dinitrogen tetroxide, oxone, molecular oxygen, (1*S*)-(+)-(10-camphorsulfonyl)oxaziridine or a peracid.

11. The method of claim 10 wherein said oxidizing reagent is iodine, *m*-chloroperbenzoic acid, iodobenzene diacetate, *tert*-butyl hydroperoxide, di-*tert*-butyl hydroperoxide, hydrogen peroxide, oxone, molecular oxygen or a peracid.

12. The method of claim 1 wherein said oxidizing reagent transfers a sulfur atom.

13. The method of claim 12 wherein said oxidizing reagent is 3-amino-1,2,4-dithiazole-5-thione; 3-ethoxy-1,2,4-dithiazoline-5-one; 1,2,4-dithiazolidine-3,5-dione; 3-methyl-1,2,4-dithiazolin-5-one; or dimethylthiuram disulfide.

5 14. The method of claim 13 wherein said oxidizing
reagent is dimethylthiuram disulfide.

15. The method of claim 1 wherein said capping reagent comprises about one part by volume of either acetic anhydride in acetonitrile or tetrahydrofuran; or chloroacetic anhydride in acetonitrile or tetrahydrofuran; added to about one part by volume of either N-methylimidazole and pyridine in acetonitrile or tetrahydrofuran; or *t*-butylphenoxyacetic anhydride in acetonitrile or tetrahydrofuran.

16. The method of claim 15 wherein said capping reagent
15 comprises about one part by volume of 20% acetic anhydride in
acetonitrile mixed with about one part by volume of a
solution having 20% N-methylimidazole, 30% pyridine and 50%
acetonitrile.

17. The method of claim 1 wherein said mixture
20 comprises dimethylthiuram disulfide, acetic anhydride,
acetoneitrile, N-methyl imidazole and pyridine.

18. The method of claim 1 wherein said mixture comprises from about 0.05M to 0.2M dimethylthiuram disulfide, about 10% acetic anhydride, about 10% N-methyl imidazole and
25 about 15% pyridine in a suitable solvent.

19. The method of claim 18 wherein said solvent is acetonitrile, toluene, ethyl acetate, tetrahydrofuran,

dichloromethane, dichloroethane, dioxane, dimethylacetamide and dimethylformamide.

20. The method of claim 1 wherein said coupling of the 5'-O-deprotected compound with the activated phosphorus composition is performed in the presence of an activating agent.

21. The method of claim 20 wherein said activating agent is 1-H-tetrazole or 4,5-dicyanoimidazole.

22. The method of claim 1 where said cyclic moiety is morpholino or phthalimido.

23. The method of claim 1 wherein each L_1 and L_2 is C_{1-6} alkyl.

24. The method of claim 23 wherein each L_1 and L_2 is isopropyl.


25. The method of claim 1 wherein L_1 and L_2 are joined together to form a heterocyclic ring system including the nitrogen atom to which said L_1 and L_2 are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S.

26. The method of claim 25 wherein said heterocyclic ring system is morpholino.


27. The method of claim 1 wherein each of said substituent groups is, independently, C_1 - C_{20} alkyl, C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, C_5 - C_{20} aryl, O-alkyl, O-alkenyl, O-alkynyl, O-aryl, O-aralkyl, O-alkylamino, O-alkylaminoalkyl (O-alkyl-N(H)alkyl), O-alkylaminodialkyl (O-alkyl-N-

(alkyl)₂), O-alkylalkoxy (O-alkyl-O-alkyl), O-alkyl-(N-imidazole), thiol, S-alkyl, S-alkenyl, S-alkynyl, NH-alkyl, NH-alkenyl, NH-alkynyl, N-dialkyl, S-aryl, NH-aryl, S-aralkyl, NH-aralkyl, N-phthalimido, halogen keto, carboxyl, 5 nitro, nitroso, nitrile, trifluoromethyl, trifluoromethoxy, N-imidazole, azido, hydrazino, hydroxylamino, isocyanato, sulfoxide, sulfone, sulfide, disulfide, silyl, heterocycle, carbocycle, polyamine, polyamide, polyalkylene glycol, or polyether;

10 or, alternatively, one or more substituent groups has one of formula I or II:



I




II

wherein:


Z₀ is O, S or NH;

15 J is a single bond, O or C(=O);

E is C₁-C₁₀ alkyl, N(R₁)(R₂), N(R₁)(R₅), N=C(R₁)(R₂), N=C(R₁)(R₅) or has one of formula III or IV;



III



IV

each R₆, R₇, R₈, R₉ and R₁₀ is, independently, hydrogen, 20 C(O)R₁₁, substituted or unsubstituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, alkylsulfonyl, arylsulfonyl, a chemical functional group or a conjugate group, wherein the substituent groups are selected from hydroxyl, amino, alkoxy,

carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl;

or optionally, R_7 and R_8 , together form a phthalimido moiety with the nitrogen atom to which they are attached;

5 or optionally, R_9 and R_{10} , together form a phthalimido moiety with the nitrogen atom to which they are attached;

each R_{11} is, independently, substituted or unsubstituted C_1 - C_{10} alkyl, trifluoromethyl, cyanoethyloxy, methoxy, ethoxy, t-butoxy, allyloxy, 9-fluorenylmethoxy, 2-(trimethylsilyl)-
10 ethoxy, 2,2,2-trichloroethoxy, benzyloxy, butyryl, iso-butyl, phenyl or aryl;

R_5 is T-L,

T is a bond or a linking moiety;

L is a chemical functional group, a conjugate group or a
15 support media;

each R_1 and R_2 is, independently, H, a nitrogen protecting group, substituted or unsubstituted C_1 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} alkenyl, substituted or unsubstituted C_2 - C_{10} alkynyl, wherein said substitution is
20 OR_3 , SR_3 , NH_3^+ , $N(R_3)(R_4)$, guanidino or acyl where said acyl is an acid amide or an ester;

or R_1 and R_2 , together, are a nitrogen protecting group or are joined in a ring structure that optionally includes an additional heteroatom selected from N and O;

25 or R_1 , T and L, together, are a chemical functional group;

each R_3 and R_4 is, independently, H, C_1 - C_{10} alkyl, a nitrogen protecting group, or R_3 and R_4 , together, are a nitrogen protecting group;

30 or R_3 and R_4 are joined in a ring structure that optionally includes an additional heteroatom selected from N and O;

Z_4 is OX, SX, or $N(X)_2$;

each X is, independently, H, C₁-C₈ alkyl, C₁-C₈ haloalkyl, C(=NH)N(H)R₅, C(=O)N(H)R₅ or OC(=O)N(H)R₅;

R₅ is H or C₁-C₈ alkyl;

Z₁, Z₂ and Z₃ comprise a ring system having from about 4
5 to about 7 carbon atoms or having from about 3 to about 6
carbon atoms and 1 or 2 hetero atoms wherein said hetero
atoms are selected from oxygen, nitrogen and sulfur and
wherein said ring system is aliphatic, unsaturated aliphatic,
aromatic, or saturated or unsaturated heterocyclic;

10 Z₅ is alkyl or haloalkyl having 1 to about 10 carbon
atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl
having 2 to about 10 carbon atoms, aryl having 6 to about 14
carbon atoms, N(R₁)(R₂)OR₁, halo, SR₁ or CN;

each q₁ is, independently, an integer from 1 to 10;

15 each q₂ is, independently, 0 or 1;

q₃ is 0 or an integer from 1 to 10;

q₄ is an integer from 1 to 10;

q₅ is from 0, 1 or 2; and

provided that when q₃ is 0, q₄ is greater than 1.

20 28. The method of claim 1 wherein said X₁ is Pg-O-, Pg-S-, CH₃-, CH₃-O-, morpholino or R₂R₃N- where each R₂ and R₃ is, independently, hydrogen or C₁-C₁₀ alkyl.

29. The method of claim 1 wherein said Pg is CH₂CH₂CN, diphenylsilylethyl, δ-cyanobutenyl, cyano p-xylyl, methyl-N-
25 trifluoroacetyl ethyl or acetoxy phenoxy ethyl.

30. The method of claim 1 wherein said heterocyclic base moiety is adenine, N⁶-benzoyladenine, cytosine, N⁴-benzoylcytosine, 5-methylcytosine, N⁴-benzoyl-5-methylcytosine, thymine, uracil, guanine, N²-isobutyrylguanine or
30 2-aminoadenine.

31. The method of claim 1 wherein said support media bound nucleoside, nucleotide, oligonucleoside or oligonucleotide is blocked at reactive sites.

5 32. The method of claim 1 wherein said blocking groups
are acid stable.

33. The method of claim 1 wherein said blocking groups are base labile.

34. The method of claim 1 wherein said deprotecting
10 reagent is acidic, neutral or basic.

35. The method of claim 32 wherein said deprotecting reagent is dichloroacetic acid, trichloroacetic acid, zinc bromide, AlCl_3 , TiCl_4 , $(\text{Et})\text{AlCl}$, $(i\text{-Bu})_2\text{AlCl}$, ceric ammonium nitrate, 1,1,1,3,3,3-hexafluoro-2-propanol or diethyloxomalonate.

36. The method of claim 35 wherein said deprotecting reagent is 2-5% dichloroacetic acid in dichloromethane or dichloroethane.

37. The method of claim 1 wherein said deprotecting
20 reagent is a fluoride moiety.

38. The method of claim 37 wherein said fluoride moiety is BF_3 -etherate.


39. The method of claim 1 wherein said oligomeric
25 compound comprises from 5 to about 50 nucleosides.

40. The method of claim 1 wherein said oligomeric compound comprises from 8 to about 30 nucleosides.

41. The method of claim 1 wherein said oligomeric compound comprises from 15 to about 25 nucleosides.

42. A method of preparing an oligomeric compound having at least one moiety of formula:

5



wherein:

X_1 is $Pg-O-$, $Pg-S-$, C_1-C_{10} straight or branched chain alkyl, $CH_3(CH_2)_{nn}-O-$, R_2R_3N- or a group remaining from coupling a chiral auxiliary;

10 nn is from 0 to 10;

Pg is CH_3 , $-CH_2CH_2CN$, $-C(CH_3)(CH_3)-CCl_3$, $-CH_2-CCl_3$, $-CH_2CH=CH_2$, $CH_2CH_2SiCH_3$, 2-yl-ethyl phenylsulfonate, δ -cyanobutenyl, cyano *p*-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-
15 trifluoroacetyl ethyl, acetoxy phenoxy ethyl, or a blocking group;


each R_2 and R_3 is, independently, hydrogen, C_1-C_{10} alkyl, cycloalkyl or aryl;

or optionally, R_2 and R_3 , together with the nitrogen
20 atom to which they are attached form a cyclic moiety that may include an additional heteroatom selected from O, S and N;

each Bx is, independently, a heterocyclic base moiety;
and

each R₁ is, independently, H, a blocked hydroxyl group, or a sugar substituent group;
comprising the steps of:

(a) providing a 5'-O-protected compound of the formula:




wherein:

T₁ is a hydroxyl protecting group; and

T₂ is a covalent attachment to a support media, or a support media bound nucleoside, nucleotide, oligonucleoside or oligonucleotide;

(b) treating said 5'-O-protected compound with a deprotecting reagent for a time and under conditions effective to form a 5'-O-deprotected compound;

(c) coupling said 5'-O-deprotected compound with an
15 activated phosphorus composition of the formula:



wherein:

T₃ is a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleoside or an oligonucleotide;

20 R_4 is $N(L_1)L_2$.

each L₁ and L₂ is, independently, C₁₋₆ straight or branched alkyl, or a C₅₋₇ cyclic aliphatic ring system;


or L₁ and L₂ are joined together to form a 4- to 13-membered heterocyclic ring system including the nitrogen atom

to which L_1 and L_2 are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S; and


R_5 is X_1 ;

- 5 or R_4 and R_5 together with the phosphorus atom to which R_4 and R_5 are attached form a chiral auxiliary;

for a time and under conditions effective to form an extended compound having the formula:



- 10 (d) treating said extended compound with dimethylthiuram disulfide in a solvent thereby forming a sulfurized compound having the formula:



(e) treating said sulfurized compound with a capping reagent for a time and under conditions effective to form said oligomeric compound.

43. The method of claim 42 further comprising treating
5 the oligomeric compound with a reagent for a time and under conditions effective to remove said blocking groups thereby forming a deblocked oligomeric compound.

44. The method of claim 43 wherein said reagent is also
10 effective to cleave the oligomeric compound from the support media.

45. The method of claim 44 wherein said reagent is aqueous ammonium hydroxide.

46. The method of claim 43 further comprising treating
15 said oligomeric compound with a further reagent for a time and under conditions effective to cleave the oligomeric compound from the support media.

47. The method of claim 42 further comprising treating
20 said oligomeric compound with a deprotecting reagent for a time and under conditions effective to deprotect the T₃ hydroxyl protecting group.

48. The method of claim 42 wherein said capping reagent
25 comprises about one part by volume of either acetic anhydride in acetonitrile or tetrahydrofuran; or chloroacetic anhydride in acetonitrile or tetrahydrofuran; added to about one part by volume of either N-methylimidazole and pyridine in acetonitrile or tetrahydrofuran; or *t*-butylphenoxyacetic anhydride in acetonitrile or tetrahydrofuran.

ISIS-4407-64204560

49. The method of claim 48 wherein said capping reagent comprises about equal volumes of 20% acetic anhydride in acetonitrile mixed with a solution having 20% N-methylimidazole, 30% pyridine and 50% acetonitrile.

5 50. The method of claim 42 wherein said solvent is acetonitrile, toluene, ethyl acetate, tetrahydrofuran, dichloromethane, dichloroethane, dioxane, dimethylacetamide and dimethylformamide.

10 51. The method of claim 42 wherein said coupling of the 5'-O-deprotected compound with the activated phosphorus composition is performed in the presence of an activating agent.

52. The method of claim 51 wherein said activating agent is 1-H-tetrazole or 4,5-dicyanoimidazole.

15 53. The method of claim 42 where said cyclic moiety is morpholino or phthalimido.

54. The method of claim 42 wherein each L_1 and L_2 is, independently, C_{1-6} alkyl.

20 55. The method of claim 54 wherein each L_1 and L_2 is isopropyl.

56. The method of claim 42 wherein L_1 and L_2 are joined together to form a heterocyclic ring system including the nitrogen atom to which said L_1 and L_2 are attached, wherein said ring system optionally includes at least one additional
25 heteroatom selected from O, N and S.

57. The method of claim 42 wherein said X_1 is Pg-O- , Pg-S- , $-\text{CH}_3$, $\text{CH}_3\text{-O-}$, morpholino or $-\text{NR}_2\text{R}_3$ where each R_2 and R_3 is, independently, hydrogen or $\text{C}_1\text{-C}_{10}$ alkyl.


58. The method of claim 42 wherein said Pg is $\text{CH}_2\text{CH}_2\text{CN}$, 5 diphenylsilylethyl, δ -cyanobutenyl, cyano *p*-xylyl, methyl-*N*-trifluoroacetyl ethyl or acetoxy phenoxy ethyl.

59. The method of claim 42 wherein said heterocyclic base moiety is adenine, N^6 -benzoyladenine, cytosine, N^4 -benzoylcytosine, 5-methylcytosine, N^4 -benzoyl-5-methyl- 10 cytosine, thymine, uracil, guanine, N^2 -isobutyrylguanine or 2-aminoadenine.

60. The method of claim 42 wherein said dimethylthiuram disulfide is from about 0.02M to about 0.2M in said solvent.

61. The method of claim 60 wherein said dimethylthiuram 15 disulfide is from about 0.1M to about 0.2M in said solvent.

62. A method of preparing an oligomeric compound having at least one moiety of one of the formulas:



wherein

20 X_2 is O or S;

X_1 is $Pg-O-$, $Pg-S-$, C_1-C_{10} straight or branched chain alkyl, $CH_3(CH_2)_{nn}-O-$, R_2R_3N- or a group remaining from coupling a chiral auxiliary;

nn is from 0 to 10;

5 Pg is CH_3 , $-CH_2CH_2CN$, $-C(CH_3)(CH_3)-CCl_3$, $-CH_2-CCl_3$, $-CH_2CH=CH_2$, $CH_2CH_2SiCH_3$, 2-yl-ethyl phenylsulfonate, δ -cyano-butenyl, cyano *p*-xylyl, diphenylsilylethyl, 4-nitro-2-yl-ethylbenzene, 2-yl-ethyl-methyl sulfonate, methyl-N-trifluoroacetyl ethyl, acetoxy phenoxy ethyl, or a blocking
10 group;


each R_2 and R_3 is, independently, hydrogen, C_1-C_{10} alkyl, cycloalkyl or aryl;

or optionally, R_2 and R_3 , together with the nitrogen atom to which they are attached form a cyclic moiety that may
15 include an additional heteroatom selected from O, S and N;

each Bx is, independently, a heterocyclic base moiety;
and

each R_1 is, independently, H, a blocked hydroxyl group, or a sugar substituent group;
20 comprising the steps of:

(a) providing a 5'-O-protected compound having one of the formulas:




or

wherein:

T_1 is a hydroxyl protecting group; and
30 T_2 is a covalent attachment to a support media, or a support media bound nucleoside, nucleotide, oligonucleoside or oligonucleotide;

(b) treating said 5'-O-protected compound with a deprotecting reagent for a time and under conditions effective to form a 5'-O-deprotected compound;

(c) coupling said 5'-O-deprotected compound with an activated phosphorus composition of the formula:



wherein:

T₃ is a hydroxyl protecting group, a nucleoside, a nucleotide, an oligonucleoside or an oligonucleotide;

10 R₄ is N(L₁)L₂;


each L₁ and L₂ is, independently, C₁₋₆ straight or branched alkyl, or a C₅₋₇ cyclic aliphatic ring system;

or L₁ and L₂ are joined together to form a 4- to 13-membered heterocyclic ring system including the nitrogen atom
 15 to which L₁ and L₂ are attached, wherein said ring system optionally includes at least one additional heteroatom selected from O, N and S; and

R₅ is X₁;

or R₄ and R₅ together with the phosphorus atom to which
 20 R₄ and R₅ are attached form a chiral auxiliary;

for a time and under conditions effective to form an extended compound having one of the formulas:



and

(d) treating said extended compound with a mixture comprising an oxidizing reagent and a capping reagent for a 5 time and under conditions effective to form said oligomeric compound.

63. A synthetic process comprising:

- adding methylamine, carbon disulfide and an organic solvent to a basic aqueous solution, thereby forming a 10 mixture;
- adding ice and acid to said mixture, thereby forming an acidified mixture;
- adding an oxidizing agent to said acidified mixture, thereby forming an oxidized mixture;
- 15 - adding a non-polar solvent to said oxidized mixture, thereby forming a precipitate;
- isolating said precipitate; and
- washing said precipitate with aqueous acid and a non-polar organic solvent.

20 64. The process of claim 63 wherein said basic aqueous solution is maintained at about 0°C during said addition of methylamine, carbon disulfide and organic solvent.

20250422 09:43:00

65. The process of claim 63 wherein said acidified mixture is maintained at about 0°C to about 5°C during said addition of said oxidizing agent.

66. The process of claim 63 wherein said basic aqueous
5 solution is aqueous sodium hydroxide.

67. The process of claim 66 wherein said sodium hydroxide has a concentration of about 2 to about 6 molar.

68. The process of claim 66 wherein the concentration of said sodium hydroxide is about 4 molar.

10 69. The process of claim 63 wherein said methylamine is added as an aqueous solution having a concentration of methylamine of about 1 to about 3M.

70. The process of claim 69 wherein said concentration of the methylamine is about 2M.

15 71. The process of claim 63 wherein said organic solvent is tetrahydrofuran.

72. The process of claim 63 wherein said acid is glacial acetic acid.

20 73. The process of claim 63 wherein said acid is added to give a final pH of about 1 to about 6.

74. The process of claim 63 wherein said oxidizing agent comprises aqueous hydrogen peroxide.

75. The process of claim 74 wherein said hydrogen peroxide has a concentration of about 10 to about 30%.

76. The process of claim 75 wherein the concentration of said hydrogen peroxide is about 30%.

77. The process of claim 63 wherein said non-polar organic solvent is hexanes or heptane.

5 78. The process of claim 63 wherein said aqueous acid
is trichloroacetic acid.

ABSTRACT OF THE DISCLOSURE

The present invention discloses methods for synthesizing oligomeric compounds. The methods include a modified
5 phosphoramidite protocol wherein the oxidation and capping steps are combined into a single step. The methods result in increased efficiency and are especially amenable to the large scale synthesis of oligomeric compounds.

DOCKET NO. ISIS-4407

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

JC862 U.S. PTO
09/640279
08/16/00

In Re Application of:

Yogesh S. Sanghvi and Quanlai Song

Group Art Unit: N/A

Examiner: N/A

For: PROCESS FOR THE PREPARATION OF
OLIGONUCLEOTIDES

DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and

I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a

☒ Utility Patent ☐ Design Patent

is sought on the invention, whose title appears above, the specification of which:

- ☒ is attached hereto.
- ☐ was filed on _____ as Serial No. _____.
- ☐ said application having been amended on _____.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to the patentability of this application in accordance with 37 CFR § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a-d) of any **foreign application(s)** for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of any application on which priority is claimed:

Priority Claimed (If X'd)	Country	Serial Number	Date Filed
<input type="checkbox"/>	_____	_____	_____
<input type="checkbox"/>	_____	_____	_____
<input type="checkbox"/>	_____	_____	_____
<input type="checkbox"/>	_____	_____	_____

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to be material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Serial Number	Date Filed	Patented/Pending/Abandoned
_____	_____	_____
_____	_____	_____
_____	_____	_____
_____	_____	_____

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below:

Serial Number	Date Filed
_____	_____
_____	_____

003130" 0201960

I hereby appoint the following persons as attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

John W. Caldwell

Registration No. 28,937

Joseph Lucci

Registration No. 33,307

of **WOODCOCK WASHBURN KURTZ MACKIEWICZ & NORRIS LLP**, One Liberty Place - 46th Floor, Philadelphia, Pennsylvania 19103 and

Herb Boswell

Registration No. 27,311

Laurel Bernstein

Registration No. 37,280

Robert S. Andrews

Registration No. 44,508

April Logan

Registration No. 33,950

of **ISIS PHARMACEUTICALS, INC.**, 2292 Faraday Avenue, Carlsbad, California 92008

Address all telephone calls and correspondence to:

Joseph Lucci, Esquire
**WOODCOCK WASHBURN KURTZ
MACKIEWICZ & NORRIS LLP**
One Liberty Place - 46th Floor
Philadelphia PA 19103
Telephone No.: **(215) 568-3100**
Facsimile No.: **(215) 568-3439**

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

00946204960

Name: Yogesh S. Sanghvi	
Mailing Address: 2169 Wandering Road Encinitas, CA 92024	Signature _____ Date of Signature: _____
City/State of Actual Residence: Encinitas, CA 92024	Citizenship: _____ <u>USA</u>

Name: Quanlai Song	
Mailing Address: 586 Barham Drive #300 San Marcos, CA 29078	Signature _____ Date of Signature: _____
City/State of Actual Residence: San Marcos, CA 29078	Citizenship: _____ <u>China</u>

20250624 15:23:53